Put your logo here



Teal. Spring. Breeding plumage. Adult. Male (18-II)

EURASIAN TEAL (Anas crecca)

IDENTIFICATION

34-38 cm. **Male** in **winter** with chesnut head with a green patch; pale breast with blackish spots; upperparts and flanks vermiculated grey; green and black speculum edged white. **Female** brown with white spotted; speculum similar to **male**.



Teal. Pattern of speculum.

SIMILAR SPECIES

Male with unmistakable plumage. The pattern of the speculum and the small size make differ the **female** from other brown ducks.

SEXING

In **breeding plumage**, **male** unmistakable with chestnut head with green band; flanks vermiculated grey; dark bill. **Female** with brown plumage; buff flanks with U-shaped pale marks; grey bill.

In **non-breeding** plumage sexes **similar** but **male** with secondaries with narrow white tips; 4th inner secondary with green speculum on all the outer vane; the innermost greater covert without pale edge; with unspotted dark bill. **Female** with secondaries with broad white tips; 4th inner secondary with green speculum only on partly of the outer vane; the innermost greater covert with pale edge; bill with dark spots.






Teal. Breeding plumage. Sexing. Pattern of head: left male; right female.



Teal. Non-breeding plumage. Sexing. Pattern of bill: left male; right female.



Teal. Sexing. Pattern of 4th secondary: left male; right female.

AGEING/SEXING

It is unlikely to find **juveniles** in Aragon, so only 2 age groups can be recognized:

1st year/2nd year spring with **juvenile** flight and tail feathers retained, which will be eroded. **Male** with the outer tertial with a quite broad black fringe on the edge; greater coverts narrow and with rounded, broad white, more or less creamy tips. **Female** with the outer tertial with a quite narrow dull black fringe on the edge; greater coverts narrow and with rounded, narrow white, more or less creamy tips.

Adult with fresh flight and tail feathers. Male with the outer tertial with a line of pearly grey, a broad line bright black and reduced pale fringe on the edge; greater coverts broad and with squared, very broad white and intensive cinnamon tips. Female with the outer tertial with a line of dark grey, a dull black line and a broad pale fringe on the edge, often around the tip; greater coverts narrow and with squared, narrow white, more or less creamy mottled grey tips.

Teal. Male. Ageing. Pattern of the innest tertial: left adult; right 1st year.



Teal. Female. Ageing. Pattern of the innest tertial: left adult; right 2nd year.





Put your logo here

Teal. Male. Ageing. Pattern of greater coverts: left adult; right 1st year.



Teal. Female. Ageing. Pattern of greater coverts: left adult; right 1st year.

MOULT

Complete **postbreeding** moult acquiring **nonbreeding** plumage, usually finished late September. Partial **postjuvenile** moult changing body feathers, tail and tertials, usually finished between October and November. Both age groups, from September, have a partial **prebre**eding moult acquiring **breeding** plumage.

PHENOLOGY

Ι	Π	III	IV	V	VI	VII	VIII	IX	Х	XI	XII

STATUS IN ARAGON


Winter visitor, with some non breeding birds remaining in summer, In winter or on passage can be found widespread throughout the Region.

Put your logo here


77 Eurasian Teal



Teal. Spring. Breeding plumage. Adult. Female (18-II)



Teal. Spring. Breeding plumage. 2nd year. Male ()



Teal. Spring. Breeding plumage. 2nd year. Head pattern: top male (); bottom female (18-II).





Teal. Spring. Breeding plumage. 2nd year. Female (18-II)





Teal. Spring. Breeding plumage. Adult. Head pattern: top male (18-II); bottom female (18-II).



Teal. Spring. Breeding plumage. Adult. Pattern of crown and bill: left male (18-II); right female (18-II).


http://blascozumeta.com

Put your logo here

77 Eurasian Teal

NO IMAGE

Teal. Spring. Breeding plumage. 2nd year. Pattern of crown and bill: left male (); right female (18-II).





Teal. Spring. Breeding plumage. 2nd year. Breast pattern: top male (); bottom female (18-II).






Teal. Spring. Breeding plumage. Adult. Breast pattern: top male (18-II); bottom female (18-II).






Teal. Spring. Breeding plumage. Adult. Upperpart pattern: left male (18-II); right female (18-II).


Put your logo here



Teal. Spring. Breeding plumage. 2nd year. Upperpart pattern: left male (); right female (18-II).



Teal. Spring. Breeding plumage. Adult. Pattern of undertail coverts: left male (18-II); right female (18-II).



Teal. Spring. Breeding plumage. 2nd year. Pattern of undertail coverts: left male (); right female (18-II).



Teal. Spring. Breeding plumage. Adult. Tail pattern: left male (18-II); right female (18-II).




Teal. Spring. Breeding plumage. 2nd year. Tail pattern: left male (); right female (18-II).



Teal. Spring. Breeding plumage. Adult. Male: pattern of flank (18-II).

Teal. Spring. Breeding plumage. Adult. Female: pattern of flank (18-II).



Teal. Spring. Breeding plumage. 2nd year. Male: pattern of flank ().





Teal. Spring. Breeding plumage. Adult. Pattern of tertials: left male (18-II); right female (18-II).




Teal. Spring. Breeding plumage. 2nd year. Pattern of tertials: left male (); right female (18-II).



Teal. Spring. Breeding plumage. 2nd year. Female: pattern of flank (18-II).




Teal. Spring. Breeding plumage. Adult. Pattern of the outermost tertial: left male (18-II); right female (18-II).



Teal. Spring. Breeding plumage. 2nd year. Pattern of the outermost tertial: left male (); right female (18-II).



Teal. Spring. Breeding plumage. Adult. Male: pattern of greater coverts (18-II).



Teal. Spring. Breeding plumage. 2nd year. Male: pattern of greater coverts ().



Teal. Spring. Breeding plumage. 2nd year. Female: pattern of greater coverts (18-II).






Teal. Spring. Breeding plumage. Adult. Female: pattern of greater coverts (18-II).

Teal. Spring. Breeding plumage. Adult. Male: pattern of primaries (18-II).



Teal. Spring. Breeding plumage. Adult. Female: pattern of primaries (18-II).


Put your logo here



Teal. Spring. Breeding plumage. 2nd year. Male: pattern of primaries ().



Teal. Spring. Breeding plumage. 2nd year. Female: pattern of primaries (18-II).



Teal. Spring. Breeding plumage. 2nd year. Pattern of the 4th inner secondary: left male (); right female (18 -II).








Teal. Spring. Breeding plumage. Adult. Pattern of the 4th inner secondary: left male (18-II); right female (18-II).

Teal. Spring. Breeding plumage. Adult. Male: pattern of secondaries (18-II).



Teal. Spring. Breeding plumage. Adult. Female: pattern of secondaries (18-II).

Put your logo here




Teal. Spring. Breeding plumage. 2nd year. Male: pattern of secondaries ().



Teal. Spring. Breeding plumage. Adult. Female: pattern of wing (18-II).



Teal. Spring. Breeding plumage. 2nd year. Female: pattern of secondaries (18-II).





Teal. Spring. Breeding plumage. Adult. Male: pattern of wing (18-II).

Teal. Spring. Breeding plumage. 2nd year. Male: pattern of wing ().



Teal. Spring. Breeding plumage. 2nd year. Female: pattern of wing (18-II).


Put your logo here



Teal. Summer. Non breeding plumage. Adult. Male (21-IX)




Teal. Summer. Non breeding plumage. 1st year. Fe-male (22-IX)



Teal. Summer. Non breeding plumage. Adult. Fe-male ()





Teal. Summer. Non breeding plumage. Adult. Head pattern: top male (21-IX); bottom female ().




Teal. Summer. Non breeding plumage. 1st year. Male (04-IX)






Teal. Summer. Non breeding plumage. 1st year. Head pattern: top male (04-IX); bottom female (22-IX).


Put your logo here



Teal. Summer. Non breeding plumage. Adult. Pattern of crown and bill: left male (21-IX); right female ().



Teal. Summer. Non breeding plumage. Adult. Breast pattern: top male (21-IX); bottom female ().



Teal. Summer. Non breeding plumage. 1st year. Pattern of crown and bill: left male (04-IX); right female (22-IX).


http://blascozumeta.com






Teal. Summer. Non breeding plumage. 1st year. Breast pattern: top male (04-IX); bottom female (22-IX).

Put your logo here



Teal. Summer. Non breeding plumage. Adult. Upperpart pattern: left male (21-IX); right female ().



Teal. Summer. Non breeding plumage. Adult. Tail pattern: left male (21-IX); right female ().




Teal. Summer. Non breeding plumage. 1st year. Upperpart pattern: left male (04-IX); right female (22-IX).




Teal. Summer. Non breeding plumage. 1st year. Tail pattern: left male (04-IX); right female (22-IX).






Teal. Summer. Non breeding plumage. Adult. Pattern of undertail coverts: left male (21-IX); right female ().



Teal. Summer. Non breeding plumage. 1st year. Pattern of undertail coverts: left male (04-IX); right female (22-IX).



Teal. Summer. Non breeding plumage. Adult. Male: pattern of flank (21-IX).



Teal. Summer. Non breeding plumage. Adult. Fe-male: pattern of flank ().




Teal. Summer. Non breeding plumage. 1st year. Male: pattern of flank (04-IX).



Teal. Summer. Non breeding plumage. 1st year. Female: pattern of flank (22-IX).






Teal. Summer. Non breeding plumage. Adult. Pattern of tertials: left male (21-IX); right female ().



Teal. Summer. Non breeding plumage. 1st year. Pattern of tertials: left male (04-IX); right female (22-IX).


Put your logo here



Teal. Summer. Non breeding plumage. Adult. Pattern of the outermost tertial: left male (21-IX); right female ().



Teal. Summer. Non breeding plumage. Adult. Male: pattern of greater coverts (21-IX).



Teal. Summer. Non breeding plumage. Adult. Fe-male: pattern of greater coverts ().



Teal. Summer. Non breeding plumage. 1st year. Pattern of the outermost tertial: left male (04-IX); right female (22-IX).




Teal. Summer. Non breeding plumage. 1st year. Male: pattern of greater coverts (04-IX).



Teal. Summer. Non breeding plumage. 1st year. Female: pattern of greater coverts (22-IX).



Teal. Summer. Non breeding plumage. Adult. Male: pattern of primaries (21-IX).




Teal. Summer. Non breeding plumage. Adult. Female: pattern of primaries ().



Teal. Summer. Non breeding plumage. 1st year. Male: pattern of primaries (04-IX).



Teal. Summer. Non breeding plumage. 1st year. Female: pattern of primaries (22-IX).



Teal. Summer. Non breeding plumage. Adult. Pattern of 4th inner secondary: left male (21-IX); right female ().




Teal. Summer. Non breeding plumage. 1st year. Pattern of 4th inner secondary: left male (04-IX); right female (22-IX).



Teal. Summer. Non breeding plumage. Adult. Male: pattern of secondaries (21-IX).


Put your logo here




Teal. Summer. Non breeding plumage. Adult. Female: pattern of secondaries ().



Teal. Summer. Non breeding plumage. Adult. Male: pattern of wing (21-IX).



Teal. Summer. Non breeding plumage. 1st year. Male: pattern of secondaries (04-IX).





Teal. Summer. Non breeding plumage. 1st year. Female: pattern of secondaries (22-IX).

Teal. Summer. Non breeding plumage. Adult. Fe-male: pattern of wing ().



Teal. Summer. Non breeding plumage. 1st year. Male: pattern of wing (04-IX).





Teal. Summer. Non breeding plumage. 1st year. Female: pattern of wing (22-IX).