

Teal. Spring. Breeding plumage. Adult. Male (18-II).

EURASIAN TEAL (Anas crecca)

IDENTIFICATION

34-38 cm. **Male** in **winter** with chesnut head with a green patch; pale breast with blackish spots; upperparts and flanks vermiculated grey; green and black speculum edged white. **Female** brown with white spotted; speculum similar to **male**.



Teal. Pattern of speculum.

SIMILAR SPECIES

Male with unmistakable plumage. The pattern of the speculum and the small size make differ the **female** from other brown ducks.

SEXING

In **breeding plumage**, **male** unmistakable with chestnut head with green band; flanks vermiculated grey; dark bill. **Female** with brown plumage; buff flanks with U-shaped pale marks; grey bill.

In **non-breeding** plumage sexes **similar** but **male** with secondaries with narrow white tips; 4th inner secondary with green speculum on all the outer vane; the innermost greater covert without pale edge; with unspotted dark bill. **Female** with secondaries with broad white tips; 4th inner secondary with green speculum only on partly of the outer vane; the innermost greater covert with pale edge; bill with dark spots.






aranzadi

Teal. Breeding plumage. Sexing. Pattern of head: left male; right female.



Teal. Non-breeding plumage. Sexing. Pattern of bill: left male; right female.





Teal. Sexing. Pattern of 4th secondary: left male; right female.

AGEING/SEXING

It is unlikely to find **juveniles** in Aragon, so only 2 age groups can be recognized:

1st year/2nd year spring with **juvenile** flight and tail feathers retained, which will be eroded. **Male** with the outer tertial with a quite broad black fringe on the edge; greater coverts narrow and with rounded, broad white, more or less creamy tips. **Female** with the outer tertial with a quite narrow dull black fringe on the edge; greater coverts narrow and with rounded, narrow white, more or less creamy tips.

Adult with fresh flight and tail feathers. Male with the outer tertial with a line of pearly grey, a broad line bright black and reduced pale fringe on the edge; greater coverts broad and with squared, very broad white and intensive cinnamon tips. Female with the outer tertial with a line of dark grey, a dull black line and a broad pale fringe on the edge, often around the tip; greater coverts narrow and with squared, narrow white, more or less creamy mottled grey tips.







Teal. Male. Ageing. Pattern of the innest tertial: left adult; right 1st year.



Teal. Female. Ageing. Pattern of the innest tertial: left adult; right 2nd year.



Teal. Male. Ageing. Pattern of greater coverts: left adult; right 1st year.



Teal. Female. Ageing. Pattern of greater coverts: left adult; right 1st year.

MOULT

Complete **postbreeding** moult acquiring **nonbreeding** plumage, usually finished late September. Partial **postjuvenile** moult changing body feathers, tail and tertials, usually finished between October and November. Both age groups, from September, have a partial **prebreeding** moult acquiring **breeding** plumage.

PHENOLOGY





Teal. Spring. Breeding plumage. Adult. Female (18-II).

PICTURE NOT AVAILABLE

PICTURE NOT AVAILABLE



Teal. Spring. B r e e d i n g plumage. 2nd year. Head pattern: top male (); bottom female (18-II).

aranzadi



Teal. Spring. Breeding plumage. 2nd year. Male ().

Teal. Spring. Breeding plumage. 2nd year. Female (18-II).





Teal. Spring. B r e e d i n g p l u m a g e . Adult. Head pattern: top male (18-II); bottom female (18-II).





Teal. Spring. Breeding plumage. Adult. Pattern of crown and bill: left male (18-II); right female (18-II).





Teal. Spring. Breeding plumage. 2nd year. Pattern of crown and bill: left male (); right female (18-II).

PICTURE NOT AVAILABLE



Teal. Spring. B r e e d i n g plumage. 2nd year. Breast pattern: top male (); bottom female (18-II).





Teal. Spring. B r e e d i n g p l u m a g e . Adult. Breast pattern: top male (18-II); bottom female (18-II).





Teal. Spring. Breeding plumage. Adult. Upperpart pattern: left male (18-II); right female (18-II).





Teal. Spring. Breeding plumage. 2nd year. Upperpart pattern: left male (); right female (18-II).



Teal. Spring. Breeding plumage. Adult. Tail pattern: left male (18-II); right female (18-II).



Teal. Spring. Breeding plumage. 2nd year. Tail pattern: left male (); right female (18-II).



Teal. Spring. Breeding plumage. Adult. Pattern of undertail coverts: left male (18-II); right female (18-II).

PICTURE NOT AVAILABLE



Teal. Spring. Breeding plumage. 2nd year. Pattern of undertail coverts: left male (); right female (18-II).



Teal. Spring. Breeding plumage. Adult. Male: pattern of flank (18-II).



Teal. Spring. Breeding plumage. Adult. Female: pattern of flank (18-II).

PICTURE NOT AVAILABLE





aranzadi zientzia elkartea

Teal. Spring. Breeding plumage. Adult. Pattern of tertials: left male (18-II); right female (18-II).





Teal. Spring. Breeding plumage. 2nd year. Pattern of tertials: left male (); right female (18-II).

Teal. Spring. Breeding plumage. 2nd year. Male: pattern of flank ().



Teal. Spring. Breeding plumage. 2nd year. Female: pattern of flank (18-II).



Teal. Spring. Breeding plumage. Adult. Pattern of the outermost tertial: left male (18-II); right female (18-II).







Teal. Spring. Breeding plumage. 2nd year. Pattern of the outermost tertial: left male (); right female (18-II).



Teal. Spring. Breeding plumage. Adult. Male: pattern of greater coverts (18-II).



Teal. Spring. Breeding plumage. Adult. Female: pattern of greater coverts (18-II).

PICTURE NOT AVAILABLE

Teal. Spring. Breeding plumage. 2nd year. Male: pattern of greater coverts ().



Teal. Spring. Breeding plumage. 2nd year. Female: pattern of greater coverts (18-II).



Teal. Spring. Breeding plumage. Adult. Male: pattern of primaries (18-II).



Teal. Spring. Breeding plumage. Adult. Female: pattern of primaries (18-II).



PICTURE NOT AVAILABLE

Teal. Spring. Breeding plumage. 2nd year. Male: pattern of primaries ().



Teal. Spring. Breeding plumage. 2nd year. Female: pattern of primaries (18-II).





Teal. Spring. Breeding plumage. 2nd year. Pattern of the 4th inner secondary: left male (); right female (18-II).





Teal. Spring. Breeding plumage. Adult. Pattern of the 4th inner secondary: left male (18-II); right female (18-II).

Teal. Spring. Breeding plumage. Adult. Male: pattern of secondaries (18-II).



Teal. Spring. Breeding plumage. Adult. Female: pattern of secondaries (18-II).



PICTURE NOT AVAILABLE

Teal. Spring. Breeding plumage. 2nd year. Male: pattern of secondaries ().



Teal. Spring. Breeding plumage. 2nd year. Female: pattern of secondaries (18-II).



Teal. Spring. Breeding plumage. Adult. Male: pattern of wing (18-II).



Teal. Spring. Breeding plumage. Adult. Female: pattern of wing (18-II).

PICTURE NOT AVAILABLE

PICTURE NOT AVAILABLE

Teal. Spring. Breeding plumage. 2nd year. Male: pattern of wing ().



Teal. Spring. Breeding plumage. 2nd year. Female: pattern of wing (18-II).





Teal. Summer. Non breeding plumage. Adult. Male (21-IX).



Teal. Summer. Non breeding plumage. 1st year. Fe-male (22-IX).



Teal. Summer. Non breeding plumage. Adult. Female ().



Teal. Summer. Non breeding plumage. 1st year. Male (04-IX).



Teal. Summer. Non b r e e d i n g p l u m a g e . Adult. Head pattern: top male (21-IX); bottom female ().








Teal. Summer. Non b r e e d i n g plumage. 1st year. Head pattern: top male (04-IX); bottom female (22-IX).






PICTURE NOT AVAILABLE

Teal. Summer. Non breeding plumage. Adult. Pattern of crown and bill: left male (21-IX); right female ().



Teal. Summer. Non breeding plumage. 1st year. Pattern of crown and bill: left male (04-IX); right female (22-IX).









Teal. Summer. Non b r e e d i n g p l u m a g e . Adult. Breast pattern: top male (21-IX); bottom female ().

Teal. Summer. Non b r e e d i n g plumage. 1st year. Breast pattern: top male (04-IX); bottom female (22-IX).






PICTURE NOT AVAILABLE

Teal. Summer. Non breeding plumage. Adult. Upperpart pattern: left male (21-IX); right female ().





Teal. Summer. Non breeding plumage. Adult. Tail pattern: left male (21-IX); right female ().




Teal. Summer. Non breeding plumage. 1st year. Upperpart pattern: left male (04-IX); right female (22-IX).




Teal. Summer. Non breeding plumage. 1st year. Tail pattern: left male (04-IX); right female (22-IX).





Teal. Summer. Non breeding plumage. Adult. Pattern of undertail coverts: left male (21-IX); right female ().





Teal. Summer. Non breeding plumage. 1st year. Pattern of undertail coverts: left male (04-IX); right female (22-IX).



Teal. Summer. Non breeding plumage. Adult. Male: pattern of flank (21-IX).



Teal. Summer. Non breeding plumage. Adult. Fe-male: pattern of flank ().




Teal. Summer. Non breeding plumage. 1st year. Male: pattern of flank (04-IX).



Teal. Summer. Non breeding plumage. 1st year. Female: pattern of flank (22-IX).





Teal. Summer. Non breeding plumage. Adult. Pattern of tertials: left male (21-IX); right female ().



Teal. Summer. Non breeding plumage. 1st year. Pattern of tertials: left male (04-IX); right female (22-IX).





PICTURE NOT AVAILABLE

Teal. Summer. Non breeding plumage. Adult. Pattern of the outermost tertial: left male (21-IX); right female ().



Teal. Summer. Non breeding plumage. 1st year. Pattern of the outermost tertial: left male (04-IX); right female (22-IX).



Teal. Summer. Non breeding plumage. Adult. Male: pattern of greater coverts (21-IX).

PICTURE NOT AVAILABLE

Teal. Summer. Non breeding plumage. Adult. Fe-male: pattern of greater coverts ().



Teal. Summer. Non breeding plumage. 1st year. Male: pattern of greater coverts (04-IX).




Teal. Summer. Non breeding plumage. 1st year. Female: pattern of greater coverts (22-IX).





Teal. Summer. Non breeding plumage. Adult. Male: pattern of primaries (21-IX).





Teal. Summer. Non breeding plumage. Adult. Pattern of 4th inner secondary: left male (21-IX); right female ().



Teal. Summer. Non breeding plumage. Adult. Female: pattern of primaries ().



Teal. Summer. Non breeding plumage. 1st year. Male: pattern of primaries (04-IX).



Teal. Summer. Non breeding plumage. 1st year. Female: pattern of primaries (22-IX).



Teal. Summer. Non breeding plumage. 1st year. Pattern of 4th inner secondary: left male (04-IX); right female (22-IX).



Teal. Summer. Non breeding plumage. Adult. Male: pattern of secondaries (21-IX).



PICTURE NOT AVAILABLE

Teal. Summer. Non breeding plumage. Adult. Female: pattern of secondaries ().



Teal. Summer. Non breeding plumage. 1st year. Male: pattern of secondaries (04-IX).



Teal. Summer. Non breeding plumage. 1st year. Female: pattern of secondaries (22-IX).





Teal. Summer. Non breeding plumage. Adult. Male: pattern of wing (21-IX).

PICTURE NOT AVAILABLE

PICTURE NOT AVAILABLE

Teal. Summer. Non breeding plumage. Adult. Fe-male: pattern of wing ().



Teal. Summer. Non breeding plumage. 1st year. Male: pattern of wing (04-IX).





Teal. Summer. Non breeding plumage. 1st year. Female: pattern of wing (22-IX).

