

Red-crested Pochard. Spring. Breeding plumage. Adult. Male ().

RED-CRESTED POCHARD (*Netta rufi-na*)

IDENTIFICATION


53-57 cm. **Male** with red-chesnut head; black breast and tail coverts; white flanks; wings chestnut with a white band; red bill. **Female** pale brown; chetnut crown and white cheeks; bill greyish with pink tip.


SIMILAR SPECIES

This species is unmistakable.


SEXING

Male in breeding plumage unmistakable with head and nape red-chestnut; breast and upper and undertail coverts black; white flanks; white leading edge of wing; reddish iris and bill; in non breeding plumage similar to female but distinguisable by red iris and bill and by white leading edge of wing. Female with brown body; head brown on crown and white on cheeks and hindneck; breast and tail coverts brownish; leading edge of wing similar as rest of wing coverts; iris reddish-brown and bill brown with pinkish tip. Juvenile birds similar but male has whiter the leading edge of wings.





Red-crested Pochard. Sexing. Pattern of head: top breeding male; middle nonbreeding male; bottom female








Red-crested Pochard. Sexing. Pattern of breast: top male; bottom female





Red-crested Pochard. Sexing. Adult: pattern of upper lesser coverts: top male; bottom female





Red-crested Pochard. Sexing. Juvenile: pattern of upper lesser coverts: top male; bottom female


AGEING

3 types of age can be recognized:

1st year summer similar to female but with narrow tail feathers with notched tips; greater coverts narrow with rounded tips; grey bill; pale brown iris.

1st year autumn/2nd year spring with juvenile flight feathers retained. In breeding plumage, male with duller and mottled underparts and narrower leading edge of wing than the adult male; female with upperparts and crown greyer-brown than the adult female.


Adult with greater coverts broad and square shaped, mainly the inner coverts; non-notched tail feathers. In breeding plumage, male with black underparts.





Red-crested Pochard. Ageing. Pattern of head and bill: top adult female; bottom juvenile





Red-crested Pochard. Ageing. Pattern of tail feathers: left female; right juvenile






Red-crested Pochard. Ageing. Pattern of greater coverts: top adult; bottom juvenile

MOULT

Complete **postbreeding** moult, acquiring eclipse plumage, usually finished in September. Partial **postjuvenile** moult changing body feathers, tail and tertials; usually finished in October. Both types of age have a partial **prebreeding** moult acquiring breeding plumage.




Red-crested Pochard. Spring. Breeding plumage. 2nd year. Male ().




Red-crested Pochard. Spring. Breeding plumage. 2nd year. Female ().

PHENOLOGY





STATUS IN ARAGON

Very scarce breeder with most of the breeding pairs in Estanca of Chiprana. More common in winter widespread thoughout the Region.





Red-crested Pochard. Spring. Breeding plumage. Adult. Female ().






Red-crested Pochard. Spring. Breeding plumage. Adult. Pattern of head and bill: top male (); bottom female ().





Red-crested Pochard. Spring. Breeding plumage. 2nd year. Pattern of head and bill: top male (); bottom female ().



Red-crested Pochard. Spring. Breeding plumage. 2nd year. Pattern of crown and bill: left male (); right female ().




http://blascozumeta.com



Red-crested Pochard. Spring. Breeding plumage. Adult. Pattern of crown and bill: left male (); right female ().






Red-crested Pochard. Spring. Breeding plumage. Adult. Breast pattern: top male (); bottom female ().




A PHOTO IS WELCOME

Red-crested Pochard. Spring. Breeding plumage. 2nd year. Breast pattern: top male (); bottom female ().



Red-crested Pochard. Spring. Breeding plumage. 2nd year. Upperpart pattern: left male (); right female ().




Red-crested Pochard. Spring. Breeding plumage. Adult. Male: pattern of flank ()



Red-crested Pochard. Spring. Breeding plumage. Adult. Female: pattern of flank ()



Red-crested Pochard. Spring. Breeding plumage. 2nd year. Male: pattern of flank ()



Red-crested Pochard. Spring. Breeding plumage. Adult. Upperpart pattern: left male (); right female ().

Put your logo here




Red-crested Pochard. Spring. Breeding plumage. 2nd year. Female: pattern of flank ()





Red-crested Pochard. Spring. Breeding plumage. 2nd year. Pattern of undertail coverts: left male (); right female ().






Red-crested Pochard. Spring. Breeding plumage. Adult. Tail pattern: left male (); right female ().






Red-crested Pochard. Spring. Breeding plumage. Adult. Pattern of tertials: left male (); right female ().





Red-crested Pochard. Spring. Breeding plumage. 2nd year. Tail pattern: left male (); right female ().






Red-crested Pochard. Spring. Breeding plumage. 2nd year. Pattern of tertials: left male (); right female ().







Red-crested Pochard. Spring. Breeding plumage. Adult. Pattern of undertail coverts: left male (); right female ().




Red-crested Pochard. Spring. Breeding plumage. Adult. Male: pattern of wing coverts ().




Red-crested Pochard. Spring. Breeding plumage. 2nd year. Female: pattern of wing coverts ().




Red-crested Pochard. Spring. Breeding plumage. Adult. Female: pattern of wing coverts ().




Red-crested Pochard. Spring. Breeding plumage. Adult. Male: pattern of primaries ()



Red-crested Pochard. Spring. Breeding plumage. 2nd year. Male: pattern of wing coverts ().




Red-crested Pochard. Spring. Breeding plumage. Adult. Female: pattern of primaries ()




Red-crested Pochard. Spring. Breeding plumage. 2nd year. Male: pattern of primaries ()




Red-crested Pochard. Spring. Breeding plumage. Adult. Female: pattern of secondaries ()




Red-crested Pochard. Spring. Breeding plumage. 2nd year Female: pattern of primaries ()



Red-crested Pochard. Spring. Breeding plumage. 2nd year. Male: pattern of secondaries ()



Red-crested Pochard. Spring. Breeding plumage. Adult. Male: pattern of secondaries ()




Red-crested Pochard. Spring. Breeding plumage. 2nd year. Female: pattern of secondaries ()




Red-crested Pochard. Spring. Breeding plumage. Adult. Male: pattern of wing ().




Red-crested Pochard. Spring. Breeding plumage. Adult. Female: pattern of wing ().




Red-crested Pochard. Spring. Breeding plumage. 2nd year. Male: pattern of wing ().




Red-crested Pochard. Spring. Breeding plumage. 2nd year. Female: pattern of wing ().




Red-crested Pochard. Summer. Non-breeding plumage. Adult. Male ().




Red-crested Pochard. Summer. Non-breeding plumage. Adult. Female ().




Red-crested Pochard. Summer. Non-breeding plumage. 2nd year. Male ().



Red-crested Pochard. Summer. Non-breeding plumage. 2nd year. Female ().







Red-crested Pochard. Summer. Nonbreeding plumage. Adult. Pattern of head and bill: top male (); bottom female ().





Red-crested Pochard. Summer. Non-breeding plumage. Adult. Pattern of crown and bill: left male (); right female ().





Red-crested
Pochard. Summer. Non-breeding plumage. 1st year.
Pattern of head and bill: top male (); bottom female ().






Red-crested Pochard. Summer. Non-breeding plumage. 1st year. Pattern of crown and bill: left male (); right female ().






Red-crested Pochard. Summer.
Non-breeding plumage. Adult.
Breast pattern: top male (); bottom female ().






Red-crested Pochard. Summer. Non-breeding plumage. Adult. Upperpart pattern: left male (); right female ().



Red-crested Pochard. Summer. Nonbreeding plumage. 1st year. Breast pattern: top male (); bottom female ().



Red-crested Pochard. Summer. Non-breeding plumage. 1st year. Upperpart pattern: left male (); right female ().






Red-crested Pochard. Summer. Non-breeding plumage. Adult. Male: pattern of flank (03-XI).




Red-crested Pochard. Summer. Non-breeding plumage. Adult. Female: pattern of flank (03-XI)





Red-crested Pochard. Summer. Non-breeding plumage. 1st year. Tail pattern: left male (); right female ().



Red-crested Pochard. Summer. Non-breeding plumage. 1st year. Male: pattern of flank ()






Red-crested Pochard. Summer. Non-breeding plumage. Adult. Pattern of undertail coverts: left male (03-XI); right female (03-XI).



Red-crested Pochard. Summer. Non-breeding plumage. 1st year. Female: pattern of flank ()








Red-crested Pochard. Summer. Non-breeding plumage. Adult. Tail pattern: left male (03-XI); right female (03-XI).

Red-crested Pochard. Summer. Non-breeding plumage. 1st year. Pattern of undertail coverts: left male (); right female ().





Red-crested Pochard. Summer. Non-breeding plumage. Adult. Pattern of tertials: left male (03-XI); right female (03-XI).






Red-crested Pochard. Summer. Non-breeding plumage. 1st year. Pattern of tertilas: left male (); right female ().




Red-crested Pochard. Summer. Non-breeding plumage. Adult. Male: pattern of wing coverts (03-XI).



Red-crested Pochard. Summer. Non-breeding plumage. Adult. Female: pattern of wing coverts (03-XI).



Red-crested Pochard. Summer. Non-breeding plumage. 1st year. Male: pattern of wing coverts ().



Red-crested Pochard. Summer. Non-breeding plumage. 1st year. Female: pattern of wing coverts ().




Red-crested Pochard. Summer. Non-breeding plumage. Adult. Male: pattern of primaries (03-XI)



Red-crested Pochard. Summer. Non-breeding plumage. Adult. Female: pattern of primaries (03-XI)




Red-crested Pochard. Summer. Non-breeding plumage. Adult. Male: pattern of secondaries (03-XI)



Red-crested Pochard. Summer. Non-breeding plumage. 1st year. Male: pattern of primaries ()



Red-crested Pochard. Summer. Non-breeding plumage. Adult. female: pattern of secondaries (03-XI)



Red-crested Pochard. Summer. Non-breeding plumage. 1st year. Female: pattern of primaries ()



Red-crested Pochard. Summer. Non-breeding plumage. 1st year. Male: pattern of secondaries ()



Red-crested Pochard. Summer. Non-breeding plumage. 1st year. Female: pattern of secondaries ()




Red-crested Pochard. Summer. Non-breeding plumage. 1st year. Female: pattern of wing ()



Red-crested Pochard. Summer. Non-breeding plumage. Adult. Male: pattern of wing (03-XI)



Red-crested Pochard. Summer. Non-breeding plumage. Adult. Female: pattern of wing (03-XI)



Red-crested Pochard. Summer. Non-breeding plumage. 1st year. Male: pattern of wing ()