

Black Kite. Spring. Adult (08-IV).

BLACK KITE (Milvus migrans)

IDENTIFICATION

59-66 cm. Brown upperparts with feathers edged pale; underparts reddish brown, streaked; pale crown with dark streaks; dark brown wings, with whitish patch on base of primaries; barred pale brown tail; black bill; yellow cere.



SIMILAR SPECIES

The only raptor with forked tail is **Red Kite** which has paler plumage tinged reddish.



SEXING

Plumage of both sexes alike. Wing length is an useful characteristic only in extreme birds: **male** usually with wing shorter than 430 mm; **female** usually with wing longer than 475 mm. **CAUTION:** overlap in wing length is so large that it is not a completely reliable characteristic for individual specimens.

AGEING

5 age groups can be recognized:

Juvenile with very fresh plumage; head and upperparts feathers with largely pale edged; upper and underwing coverts with pale edge; greater and primary coverts with pale tips forming a thin line across the wing; breast feathers light brown with a dark shaft streak; dark brown iris.

2nd year spring with juvenile plumage but

2nd year spring with juvenile plumage but less contrasted due to wear; body with some adult-type feathers; dark brown iris.

2nd year autumn with two ages of feathers, with brownish and worn **juvenile** feathers contrasting with darker and fresh moulted ones; dark eyes.

3rd year only in birds retaining some unmoulted **juvenile** flight feathers, which are very old and worn, or **2nd year** feathers, shorter and pointed than **adult** type; iris light brown.

Adult with greyish head with thin dark streaks; neck and upperparts with grey brown feathers without pale edges; upper and underwing coverts without pale edge; greater and primary coverts without pale tips forming a line; breast feathers dark brown with a dark central streak; pale iris. CAUTION: it is necessary more information about when adults have pale iris and how important is individual variation.





Black Kite. Ageing. Pattern of head and iris colour: left adult; right juvenile.






Black Kite. Ageing. Pattern of breast: left adult; right juvenile.

PHENOLOGY

MOULT





STATUS IN ARAGON

Summer visitor. Widely distributed throughout the Region, more scarce in Southern areas.



Black Kite. Ageing. Pattern of wing coverts: top adult; bottom juvenile.



Black Kite. Spring. 2nd year (13-V).





Black Kite. Ageing. Pattern of primary coverts: top adult; bottom juvenile.



Black Kite. Spring. 3rd year (08-IV).





Black Kite. Spring. Head pattern and iris colour: top adult (08 -IV); middle 3rd year (08-IV); bottom 2nd year (13-V).







Black Kite. Spring. Breast pattern: top left adult (08-IV); top right 3rd year (08-IV); left 2nd year (13-V).









Black Kite. Spring. Crown pattern: top left adult (08-IV); top right 3rd year (08-IV); left 2nd year (13-V).







Black Kite. Spring. Upperpart pattern: top left adult (08-IV); top right 3rd year (08-IV); left 2nd year (13-V).





Black Kite. Spring.
Tail pattern: top adult (08-IV); middle 3rd year (08-IV); bottom 2nd year (19-V).





Black Kite. Spring. Adult: pattern of primary coverts (08-IV).



Black Kite. Spring. 3rd year: pattern of primary coverts (08-IV).



Black Kite. Spring. 2nd year: pattern of primary coverts 19-V).



Black Kite. Spring. Adult: pattern of primaries (08-IV).



Black Kite. Spring. 3rd year: pattern of primaries (08 -IV).



Black Kite. Spring. 2nd year: pattern of primaries (19-V).



Black Kite. Spring. Adult: pattern of wing coverts and secondaries (08-IV).



Black Kite. Spring. 3rd year: pattern of wing coverts and secondaries (08-IV).



Black Kite. Spring. 2nd year: pattern of wing coverts and secondaries (19-V).



Black Kite. Spring. Adult: pattern of underwing coverts (08-IV).



Black Kite. Spring. 3rd year: pattern of underwing coverts (08-IV).



Black Kite. Spring. 3rd year: pattern of wing (1 2nd year feathers) (08-IV).



Black Kite. Spring. 2nd year: pattern of underwing coverts (19-V).





Black Kite. Spring. 2nd year: pattern of wing (19-V).





Black Kite. Spring. Adult: pattern of wing (08-IV).



Black Kite. Summer. Adult (05-VII).



Black Kite. Summer. 3rd year (22-VII)



Black Kite. Summer. 2nd year (13-VII).



Black Kite. Summer. Juvenile (25-VII).



http://blascozumeta.com

Black Kite. Summer. Head pattern and iris colour: top adult (05-VII); bottom 3rd year (22-VII)





Black Kite. Summer. Head pattern and iris colour: top 2nd year (13-VII); bottom juvenile (25-VII).





Black Kite. Summer. Crown pattern: left adult (05-VII); right 3rd year (22-VII)





Black Kite. Summer. Crown pattern: left 2nd year (22-VII); right juvenile (25-VII).





Black Kite. Summer. Breast pattern: left adult (05-VII); right 3rd year (22-VII).





Black Kite. Summer. Breast pattern: left 2nd year (13-VII); right juvenile (25-VII).





Black Kite. Summer. Upperpart pattern: left adult (05-VII); right 3rd year (22-VII)





Black Kite. Summer. Upperpart pattern: left 2nd year (13-VII); right juvenile (25-VII).





Black Kite. Summer. Tail pattern: left adult (05-VII); right 3rd year (22-VII).





Black Kite. Summer. Tail pattern: left 2nd year (13-VII); right juvenile (25-VII).



Black Kite. Summer. Adult: pattern of primary coverts (03-VII).



Black Kite. Summer. Juvenile: pattern of primary coverts (25-VII).



Black Kite. Summer. 3rd year: pattern of primary coverts (22-VII)



Black Kite. Summer. Adult: pattern of primaries (03-VII).



Black Kite. Summer. 2nd year: pattern of primary coverts (13-VII).



Black Kite. Summer. 3rd year: pattern of primaries (22-VII)



Black Kite. Summer. 2nd year: pattern of primaries (13-VII).



Black Kite. Summer. Juvenile: pattern of primaries (25-VII).



Black Kite. Summer. Adult: pattern of wing coverts and secondaries (03-VII).



Black Kite. Summer. 3rd year: pattern of wing coverts and secondaries (1 juvenile feather; 2 2nd year feather; 3 3rd year feather) (22-VII)



Black Kite. Summer. 2nd year: pattern of wing coverts and secondaries (13-VII).



Black Kite. Summer. Juvenile: pattern of wing coverts and secondaries (25-VII).



Black Kite. Summer. Adult: pattern of underwing coverts (03-VII).





Black Kite. Summer. Adult: pattern of wing (03-VI).



Black Kite. Summer. 3rd year: pattern of underwing coverts (22-VII)





Black Kite. Summer. 3rd year: pattern of wing (1 juvenile feather; 2 2nd year feather; 3 3rd year feather) (22-VII)



Black Kite. Summer. 2nd year: pattern of underwing coverts (13-VII).



Black Kite. Summer. Juvenile: patrern of underwing coverts (25-VII)





Black Kite. Summer. 2nd year: pattern of wing (13-VII).





Black Kite. Summer. Juvenile: pattern of wing (25-VII)