

Hen Harrier. Adult. Male (08-III)

Hen Harrier. Adult. Male. Pattern of wing.

#### **NORTHERN HARRIER** (Circus cyaneus)


#### **IDENTIFICATION**

43-51 cm. Adult male with grey head and upperparts; underparts grey on breast and white on vent; black wing-tips; grey tail; white rump. Adult female with dark brown upperparts; underparts light brown with dark streaked; white rump; dark barred tail and wings. Juvenile similar to females, but with underparts tinged rusty-yellow. Length of 10th primary between 5th and 6th; the beginning of emargination of 9th primary hidden by primary coverts.

#### SIMILAR SPECIES

Adult male similar to male of Montagu's Harrier, with smaller white rump, with a dark wing bar and streakes flanks and vent. Females of Montagu's Harrier have smaller white rump and have a band just below the wing coverts; 10th primary with similar length than 6th; the beginning of emargination of 9th primary is not hiden by primary coverts. Female and juveniles of Marsh Harrier have a pale patch on head and throat.





Hen Harrier. Adult. Female. Wing formula: emargination of 9th primary hidden by primary coverts; length of 10th primary between 5th and 6th.







#### AGEING/SEXING

3 age groups can be recognized:

Juvenile with one age of very fresh feathers; head well contrasted with dark cheeks and crown; upperparts with dark brown feathers edged rufous; ochre underparts with dark streaks; tip of greater and primary coverts forming a clear thin line; outermost tail feather with well-defined subterminal dark band; juveniles cannot be sexed by plumage pattern but iris colour and size are useful characteristics: male with pale yellow, grey or grey-brown iris; wing shorter than 340 mm; female with dark brown iris; wing longer than 355 mm.

2nd year autumn/3rd year spring male with juvenile feathers often retained on crown, neck or breast and sometimes some greater coverts; mantle and scapulars tinged dark brownish-grey; usually central tail feathers with ill-defined dark bands; outermost tail feather with well-defined dark line on the outer web.

2nd year autumn/3rd year spring female similar to adult female when moult has been completed; iris brown or golden brown; outermost tail feather with well-defined subterminal dark band.

Adult male with bluish-grey, white and black plumage; yellow iris; without brown feathers on head and wings; blue-grey upperparts without brownish tinge; white underwing coverts; if some flight feathers remain unmoulted then with adult pattern; usually central tail feathers without dark bands; outermost tail feather with ill-defined dark line on the outer web.


Adult female with yellow iris (CAUTION: the bright yellow iris is acquired at 5-7 years old); head weakly contrasted with an extended periocular white area; dark brown upperparts; underparts whitish with brown streaks; from below, flight feathers whitish banded black; underwing coverts mottled buff; greater coverts without pale tips; outermost tail feather with ill-defined subterminal dark band.





Hen Harrier. Adult. Sexing. Pattern of breast and head: left male; right female.









Hen Harrier. Juvenile. Sexing. Colour of iris: top male; bottom female.



Hen Harrier.
Ageing.
Pattern of greater coverts:
top adult female;
bottom
juvenile.





Hen Harrier. Male. Ageing. Pattern of head: top adult; bottom 2nd year.







Hen Harrier. Female. Ageing. Pattern of the outermost tail feather: left adult; middle 2nd year; right juvenile.





Hen Harrier. Male. Ageing. Pattern of rump: left adult; right 2nd year.



Hen Harrier. Male. Ageing. Pattern of nape: left adult; right 2nd year.

### **MOULT**

Complete **postbreeding** moult, starting in April and finished in November; some birds may have unmoulted secondaries. Partial **postjuvenile** moult reduced to some body feathers changed during the winter; the first complete moult takes place in the **2nd year** at the same period as **adults**; occasionally some birds may have unmoulted body and flight feathers.

#### **PHENOLOGY**



### **STATUS IN ARAGON**

Mainly a winter visitor, with a little pairs of breeding birds in the Ebro Valley, Gallocanta and South of Teruel; in winter a common species in plain cultivated areas of the region.



Hen Harrier. Adult. Female (30-I)



Hen Harrier. Juvenile. Male (14-II)



Hen Harrier. Juvenile. Female (30-I)



Hen Harrier. 2nd year autumn. Male (25-XII)



Hen Harrier. Adult. Head pattern: top male (08-III); bottom female (30-I).



Hen Harrier. 2nd year autumn. Female (25-XII)







Hen Harrier. 2nd year autumn. Head pattern: top male with all feathers moulted (25-XII); middle male with some retained juvenile feathers (25-XII); bottom female (25-XII).







Hen Harrier. Adult. Crown pattern: left male (08-III); right female (30-I).





Hen Harrier. 2nd year autumn. Crown pattern: left male (25-XII); right female (25-XII).





Hen Harrier. Juvenile. Head pattern: top male (14-II); bottom female (30-I).





Hen Harrier. Juvenile. Crown pattern: left male (14-II); right female (30-I).






Hen Harrier. Adult. Nape pattern: left male (08-III); right female (30-I).





Hen Harrier. 2nd year autumn. Nape pattern: left male (25-XII); right female (25-XII).







Hen Harrier. Juvenile. Breast pattern: left male (14-II); right female (30-I).





Hen Harrier. Juvenile. Nape pattern: left male (14-II); right female (30-I).



Hen Harrier. Adult. Flank pattern: top male (08-III); bottom female (30-I).





Hen Harrier. Adult. Breast pattern: left male (08-III); right female (30-I).







Hen Harrier. 2nd year autumn. Flank pattern: top male (25-XII); bottom female (25-XII).







Hen Harrier. 2nd year autumn. Breast pattern: top left male with all feathers moulted (25-XII); top right male with some retained juvenile feathers (25-XII); left female (25-XII).



Hen Harrier.
Juvenile.
Flank
pattern:
top male
(14-II);
bottom
female
(30-I).







Hen Harrier. Juvenile. Upperpart pattern: left male (14-II); right female (30-I).





Hen Harrier. Adult. Upperpart pattern: left male (08-III); right female (30-I).

Hen Harrier. Adult. Rump pattern: left male (08-III); right female (30-I).







Hen Harrier. 2nd year autumn. Upperpart pattern: left male (25-XII); right female (25-XII).

Hen Harrier. 2nd year autumn. Rump pattern: top left male with all feathers moulted (25-XII); top right male with some retained juvenile feathers (25-XII); left female (25-XII).







Hen Harrier. Juvenile. Rump pattern: top left male with moulted feathers (14-II); top right male with unmoulted feathers (14-II); left female (30-I).





Hen Harrier. 2nd year autumn. Male: tail pattern (25-XII).





Hen Harrier. 2nd year autumn. Female: tail pattern (25-XII).





Hen Harrier. Adult. Male: tail pattern (08-III)





Hen Harrier. Juvenile. Male: tail pattern (14-II).





Hen Harrier. Adult. Female: tail pattern (30-I)





Hen Harrier. Juvenile. Female: tail pattern (30-I).



Hen Harrier.
Adult.
Pattern of primary coverts: top male (08-III); bottom female (30-I).





Hen Harrier.
Adult.
Pattern of underwing coverts: top male (08-III); bottom female (30-I).





Hen Ha-2nd rrier. year autumn. Pattern of primary coverts: top male (25-XII); bottom female (1 adult feather; 2 juvenile feather)

(25-XIÍ).





Hen Harrier. 2nd year autumn. Pattern of underwing coverts: top male (25-XII); bottom female (25-XII).





Hen Harrier. Juvenile. Pattern of primary coverts: top male (14-II); bottom female (30-I).





Hen Harrier. Juvenile. Pattern of underwing coverts: top male (14-II); bottom female (30-I).





Hen Harrier. Adult. Male: pattern of primaries (08-III).





Hen Harrier. 2nd year autumn. Male: pattern of primaries (25-XII).





Hen Harrier. Adult. Female: pattern of primaries (30-1)





Hen Harrier. 2nd year autumn. Female: pattern of primaries (1 adult feathers; 2 juvenile feathers) (25-XII).





Hen Harrier. Juvenile. Male: pattern of primaries (14 -II).





Hen Harrier. Adult. Male: pattern of secondaries(08-III).





Hen Harrier. Juvenile. Female: pattern of primaries (30-I).





Hen Harrier. Adult. Female: pattern of secondaries (30-I).





Hen Harrier. 2nd year autumn. Male: pattern of secondaries and wing coverts with all feathers moulted (25-XII).





Hen Harrier. 2nd year autumn. Female: pattern of secondaries (1 adult feathers; 2 juvenile feathers) (25 -XII).





Hen Harrier. 2nd year autumn. Male: pattern of secondaries and wing coverts with juvenile feathers retained (25-XII).





Hen Harrier. Juvenile. Male: pattern of secondaries (14-II).





Hen Harrier. Juvenile. Female: pattern of secondaries (30-1).





Hen Harrier. Adult. Male: pattern of wing (08-III).





Hen Harrier. Adult. Female: pattern of wing (30-I).





Hen Harrier. 2nd year autumn. Male: pattern of wing with all feathers moulted (25-XII).





Hen Harrier. 2nd year autumn. Male: pattern of wing with juvenile feathers retained (25-XII).





Hen Harrier. 2nd year autumn. Female: pattern of wing (1 juvenile feathers; 2 adult feathers) (25-XII).





Hen Harrier. Juvenile. Female: pattern of wing (30-I).



Hen Harrier. Juvenile. Male: pattern of wing (14-II).