

Spotted Crake. Spring. Adult. Male (21-III)

SPOTTED CRAKE (*Porzana porzana*)

IDENTIFICATION

21-23 cm. Head with black spotted brown crown and grey band on sides; feathers on upperparts and wing coverts dark brown, with olive edges and white streaks; breast white spotted; underparts grey with olive tinge; flanks barred with brown, black and white bars; greenyellowish bill with red base; green olive legs.





Spotted Crake. Pattern of head and undertail coverts.




SIMILAR SPECIES

Water Rail has a long and red bill. Little Crake lacks white speckling on head, breast and nape. Baillon's Crake lacks white speckling on head and without red spot on base of bill.







SEXING


In **spring**, during **breeding** period, **male** with supercilium, cheeks and throat pure slate grey with only a small area of white speckling bordering ear coverts; most **males** with bright and contrasted bill; wing longer than 124 mm. **Female** with supercilium, cheeks and throat dull slate grey with heavy white speckling bordering ear coverts; some **females** with dull and slightly contrasted bill; wing shorter than 116 mm.

In autumn, adult male with cheeks and throat heavily speckled with white; supercilium grey with some white speckling mainly to the rear of head. Adult female similar to male, but throat and cheeks heavily speckled white with pure grey restricted to front part of supercilium and to central chin (CAUTION: in both seasons, sexes have great overlap in pattern, colours and biometrics, so these characteristics are useful only for extreme birds).

1st year birds cannot be sexed using plumage pattern.



Spotted Crake. Spring. xing. Pattern of head: top male; bottom female







Spotted Crake. Spring. Ageing. Wear of primaries: top adult; bottom 2nd year.



Spotted Crake. Autumn. Sexing. Pattern of head: top male; bottom female



A PHOTO IS WELCOME

Spotted Crake. Ageing. Pattern of head and colour of bill: top adult; bottom 1st year.


AGEING

This species is unlikely to breed in Aragon, so only 3 age groups are described:

1st year with withish throat; sides of head and supercilium with extensive white speckles; dark brown iris.

2nd year spring similar to adults and many indistinguishable, but can retain old pale juvenile feathers on chin and throat; with juvenile flight feathers.

Adult with plumage described in SEXING; pale brown iris.





Spotted Crake. Ageing. Pattern of chin: left adult; right 1st year.

MOULT

Complete **postbreeding** moult, usually finished in October. Partial **postjuvenile** moult confined to body feathers; generally finished in October. Both age classes have a **prebreeding** moult confined to body feathers.

PHENOLOGY


I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XII

STATUS IN ARAGON

Poorly known status. There are some records in breeding period, being a more common species on passage.



Spotted Crake. Spring. Adult. Female (21-III)



Spotted Crake. Spring. 2nd year. Male ()



Spotted Crake. Spring. 2nd year. Female (21-III)




Spotted Crake. Spring. Adult. Head pattern: top male (21-III); bottom female (21-III).







Spotted Crake. Spring. 2nd year. Head pattern: top male (25-IV) (Photo: Reinhard Vohwinkel); bottom female (21-III).





Spotted Crake. Spring. Adult. Crown pattern: left male (21-III); right female (21-III).





Spotted Crake. Spring. 2nd year. Chin pattern: left male (); right female (21-III).





Spotted Crake. Spring. 2nd year. Crown pattern: left male (Photo: Reinhard Vohwinkel); right female (21-III).





Spotted Crake. Spring. Adult. Breast pattern: left male (21-III); right female (21-III).





Spotted Crake. Spring. Adult. Chin pattern: left male (21-III); right female (21-III).






Spotted Crake. Spring. 2nd year. Breast pattern: left male (Photo: Reinhard Vohwinkel); right female (21-III).





Spotted Crake. Spring. Adult. Pattern of upperparts: left male (21-III); right female (21-III).





Spotted Crake. Spring. 2nd year. Pattern of legs: top male (); bottom female (21-III).





Spotted Crake. Spring. 2nd year. Pattern of upperparts: left male (); right female (21-III).





Spotted Crake. Spring. Adult. Tail pattern: left male (21-III); right female (21-III).





Spotted Crake. Spring. Adult. Pattern of legs: top male (21-III); bottom female (21-III).





Spotted Crake. Spring. 2nd year. Tail pattern: left male (Photo: Reinhard Vohwinkel); right female (21-III).





Spotted Crake. Spring. Adult. Pattern of undertail coverts: left male (21-III); right female (21-III).





Spotted Crake. Spring. 2nd year. Pattern of undertail coverts: left male (Photo: Reinhard Vohwinkel); right female (21-III).



Spotted Crake. Spring. Adult. Male: pattern of flank (21-III).



Spotted Crake. Spring. Adult. Female: pattern of flank (21-III).



Spotted Crake. Spring. 2nd year. Male: pattern of flank (Photo: Reinhard Vohwinkel).



Spotted Crake. Spring. 2nd year. Female: pattern of flank (21-III).





Spotted Crake. Spring. Adult. Pattern of tertials: left male (21-III); right female (21-III).





Spotted Crake. Spring. 2nd year. Pattern of tertials: left male (Photo: Reinhard Vohwinkel); right female (21-III).



Spotted Crake. Spring. Adult. Male: pattern of primary coverts (21-III).



Spotted Crake. Spring. Adult. Female: pattern of primary coverts (21-III).



Spotted Crake. Spring. 2nd year. Male: pattern of primary coverts (Photo: Reinhard Vohwinkel).




Spotted Crake. Spring. 2nd year. Female: pattern of primary coverts (21-III).



Spotted Crake. Spring. Adult. Male: pattern of primaries (21-III).



Spotted Crake. Spring. Adult. Female: pattern of primaries (21-III).



Spotted Crake. Spring. 2nd year. Male: pattern of primaries ().



Spotted Crake. Spring. 2nd year. Female: pattern of primaries (21-III).



Spotted Crake. Spring. Adult. Male: pattern of secondaries (21-III).



Spotted Crake. Spring. Adult. Female: pattern of secondaries (21-III).



Spotted Crake. Spring. 2nd year. Male: pattern of secondaries (Photo: Reinhard Vohwinkel).



Spotted Crake. Spring. 2nd year. Female: pattern of secondaries (21-III).






Spotted Crake. Spring. Adult. Male: pattern of wing (21-III).





Spotted Crake. Spring. Adult. Female: pattern of wing (21-III).






Spotted Crake. Spring. 2nd year. Male: pattern of wing ().





Spotted Crake. Spring. 2nd year. Female: pattern of wing (21-III).




Spotted Crake. Autumn. Adult. Male ().





Spotted Crake. Autumn. Head pattern and bill colour: top adult male (); middle adult female (); bottom 1st year (28-X).



Spotted Crake. Autumn. Adult. Female ().











Spotted Crake. Autumn. 1st year (28-X).



Spotted Crake. Autumn. Crown pattern: top left adult male (); top right adult female (); left 1st year (28-X).











Spotted Crake. Autumn. Chin pattern: top left adult male (); top right adult female (); left 1st year (28-X).




Spotted Crake. Autumn. Upperpart pattern: top left adult male (); top right adult female (); left 1st year (28-X).







Spotted Crake. Autumn. Breast pattern: top left adult male (); top right adult female (); left 1st year (28-X).








Spotted Crake. Autumn. Pattern fo legs: top adult male (); middle adult female (); bottom 1st year (28-X).








Spotted Crake. Autumn. Tail pattern: top left adult male (); top right adult female (); left 1st year (28-X).




Spotted Crake. Autumn. Adult. Male: pattern of flank ().



Spotted Crake. Autumn. Adult. Female: pattern of flank ().







Spotted Crake. Autumn. Pattern of undertail coverts: top left adult male (); top right adult female (); left 1st year (28-X).




Spotted Crake. Autumn. 1st year: pattern of flank (28 -X)








Spotted Crake. Autumn. Pattern of tertials: top left adult male (); top right adult female (); left 1st year (28-X).




Spotted Crake. Autumn. Adult. Female: pattern of primary coverts ().




Spotted Crake. Autumn. 1st year: pattern of primary coverts (28-X).



Spotted Crake. Autumn. Adult. Male: pattern of primary coverts ().




Spotted Crake. Autumn. Adult. Male: pattern of primaries ().




Spotted Crake. Autumn. Adult. Female: pattern of primaries ().



Spotted Crake. Autumn. 1st year: pattern of primaries (28-X).




Spotted Crake. Autumn. Adult. Male: pattern of secondaries ().



Spotted Crake. Autumn. Adult. Female: pattern of secondaries ().



Spotted Crake. Autumn. 1st year: pattern of secondaries (28-X).






Spotted Crake. Autumn. Adult. Male: pattern of wing ().





Spotted Crake. Autumn. 1st year: pattern of wing (28 -X).





Spotted Crake. Autumn. Adult. Female: pattern of wing ().