




Black-winged Stilt. Adult. Male (22-IV)

BLACK-WINGED STILT (Himantopus himantopus)

IDENTIFICATION

37-38 cm. White plumage except dark upperparts; usually black on head and neck; black bill; red and very long legs.



SIMILAR SPECIES

This species is unmistakable.

SEXING

In adult birds, male with upperparts, scapulars and tertials black; wing longer than 240 mm. Female with upperparts, scapulars and tertials dark brown, contrasting with black wing coverts; wing shorter than 235 mm (CAUTION: in breeding plumaje pattern of black colour on head is strongly variable in both sexes; in non breeding plumage both sexes have similar grey pattern). Sexing of juveniles recorded by some authors (male with wing coverts with metallic green gloss; female with wing coverts without metallic green gloss) doubtful due to overlap and must be verified.



Black-winged Stilt. Adult. Sexing. Pattern of upperparts: left male; right female.

AGEING

3 age groups can be recognized:

Juvenile with fresh plumage; feathers of upperparts, scapulars, tertials and wing coverts clearly edged buff; secondaries and inner primaries with white tips; crown greyish, never white; pale pink legs.

2nd year similar to **adult female** but with **juvenile** flight feathers, which are worn and have white tips; sometimes with **juvenile** tertials and wing coverts retained.

Adult with upperparts, scapulars, tertials and wing coverts plain black; secondaries and inner primaries without pale tips; crown white or with a black patch; deep pink legs.





Black-winged Stilt. Ageing. Pattern of head: top adult; bottom juvenile.





Black-winged Stilt. Ageing. Pattern of wing coverts: top adult; bottom juvenile.







Black-winged Stilt. Ageing. Pattern of upperparts: left adult; right juvenile.




Black-winged Stilt. Ageing. Colour of legs: top adult; bottom juvenile.



MOULT

Complete **postbreeding** moult, suspended during migration and finished in wintering sites. Partial **postjuvenile** moult involving body feathers, some central tail feathers and all or most of lesser and median wing coverts; starts in July and finished in winter quarters. Both age classes have a partial **prebreeding** moult including body feathers, some scapulars, tertials and central tail feathers.

PHENOLOGY





Black-winged Stilt. Adult. Female (22-IV)

http://blascozumeta.com http://www.aranzadi.eus ²



PICTURE NOT AVAILABLE

Black-winged Stilt. 2nd year. Male ()

PICTURE NOT AVAILABLE

Black-winged Stilt. 2nd year. Female ()



Black-winged Stilt. Juvenile (21-VII).



Black-winged Stilt. Young (22-VI)





Black-winged Stilt. Adult. Head pattern: top male (22-IV); bottom female (22-IV)



PICTURE NOT AVAILABLE

PICTURE NOT AVAILABLE

Black-winged Stilt. Head pattern: top 2nd year male (); middle 2nd year female (); bottom juvenile (21-VII).



PICTURE NOT AVAILABLE



Black-winged Stilt. Nape pattern: top left 2nd year male (); top right 2nd year female (); left juvenile (21-VII).







Black-winged Stilt. Adult. Nape pattern: left male (22-IV); right female (22-IV)





Black-winged Stilt. Adult. Crown pattern: left male (22-IV); right female (22-IV)



PICTURE NOT AVAILABLE

PICTURE NOT AVAILABLE



Black-winged Stilt. Crown pattern: top left 2nd year male (); top right 2nd year female (); left juvenile (21-VII).



Blackwinged Stilt. Adult. Breast pattern: top male (22-IV); bottom female (22-IV)



PICTURE NOT AVAILABLE

PICTURE NOT AVAILABLE

Black-winged Stilt. Head pattern: top 2nd year male (); middle 2nd year female (); bottom juvenile (21-VII).







5

Black-winged Stilt. Adult. Upperpart pattern: left male (22-IV); right female (22-IV)



PICTURE NOT AVAILABLE

PICTURE NOT AVAILABLE

PICTURE NOT AVAILABLE

PICTURE NOT AVAILABLE

Black-winged Stilt. Legs pattern: top 2nd year male (); middle 2nd year female bottom juvenile (21-VII).




Black-winged Upperpart pattern: top left 2nd year male (); top right 2nd year female (); left juvenile (21-VII).



Black-winged Stilt. Adult. Legs pattern: top male (22-IV); bottom female (22-IV)







Black-winged Stilt. Adult. Tail pattern: left male (22-IV);

right female (22-IV)



PICTURE NOT AVAILABLE

PICTURE NOT AVAILABLE

PICTURE NOT AVAILABLE

PICTURE NOT AVAILABLE



Black-winged Stilt. Tail pattern: top left 2nd year male (); top right 2nd year female (); left juvenile (21-VII).



Black-winged Stilt. Pattern of tertials: top left 2nd year male (); top right 2nd year female (); left juvenile (21-VII).







Black-winged Stilt. Adult. Pattern of tertials: left male (22-IV); right female (22-IV)

Black-winged Stilt. Adult. Male: pattern of wing coverts (22 -IV).





Black-winged Stilt. Adult. Female: pattern of wing coverts (22-IV).



Black-winged Stilt. Juvenile: pattern of wing coverts (21-

VII)

Black-winged Stilt. Adult. Male: pattern of primaries (22-IV)

PICTURE NOT AVAILABLE

Black-winged Stilt. 2nd year. Male: pattern of wing coverts ().



Black-winged Stilt. Adult. Female: pattern of primaries (22-IV)

PICTURE NOT AVAILABLE

Black-winged Stilt. 2nd year. Female: pattern of wing coverts ().

PICTURE NOT AVAILABLE

Black-winged Stilt. 2nd year. Male: pattern of primaries ()



PICTURE NOT AVAILABLE

Black-winged Stilt. 2nd year. Female: pattern of primaries ()



Black-winged Stilt. Juvenile: pattern of primaries (21-VII)



Black-winged Stilt. Adult. Male: pattern of secondaries (22-IV)



Black-winged Stilt. Adult. Female: pattern of secondaries (22-IV)

PICTURE NOT AVAILABLE

Black-winged Stilt. 2nd year. Male: pattern of secondaries

PICTURE NOT AVAILABLE

Black-winged Stilt. 2nd year. Female: pattern of secondaries ()





Black-winged Stilt. Juvenile: pattern of secondaries (21-VII)



Black-winged Stilt. Adult. Male: pattern of wing (22-IV)





Black-winged Stilt. Adult. Female: pattern of wing (22-IV)



PICTURE NOT AVAILABLE

Black-winged Stilt. 2nd year. Male: pattern of wing ()

PICTURE NOT AVAILABLE

PICTURE NOT AVAILABLE

Black-winged Stilt. 2nd year. Female: pattern of wing ()





Black-winged Stilt. Juvenile: pattern of wing (21-VII)