Put your logo here



Black-winged Stilt. Adult. Male (22-IV)

BLACK-WINGED STILT (Himantopus *himantopus*)

IDENTIFICATION

37-38 cm. White plumage except dark upperparts; usually black on head and neck; black bill; red and very long legs.



Black-winged Stilt. Pattern of head an legs.



SIMILAR SPECIES

This species is unmistakable.

SEXING

In **adult** birds, **male** with upperparts, scapulars and tertials black; wing longer than 240 mm. Female with upperparts, scapulars and tertials dark brown, contrasting with black wing coverts; wing shorter than 235 mm (CAUTION: in breeding plumaje pattern of black colour on head is strongly variable in both sexes; in non breeding plumage both sexes have similar grey pattern). Sexing of juveniles recorded by some authors (male with wing coverts with metallic green gloss; female with wing coverts without metallic green gloss) doubtful due to overlap and must be verified.



Black-winged Stilt. Adult. Sexing. Pattern of upperparts: left male; right female.

AGEING

3 age groups can be recognized:

Juvenile with fresh plumage; feathers of upperparts, scapulars, tertials and wing coverts clearly edged buff; secondaries and inner primaries with white tips; crown greyish, never white; pale pink legs.

2nd year similar to adult female but with juvenile flight feathers, which are worn and have white tips; sometimes with juvenile tertials and wing coverts retained.

Adult with upperparts, scapulars, tertials and wing coverts plain black; secondaries and inner primaries without pale tips; crown white or with a black patch; deep pink legs.

Put your logo here





Black-





Black-winged Stilt. Ageing. Pattern of wing coverts: top adult; bottom juvenile.

MOULT

Complete **postbreeding** moult, suspended during migration and finished in wintering sites. Partial **postjuvenile** moult involving body feathers, some central tail feathers and all or most of lesser and median wing coverts; starts in July and finished in winter quarters. Both age classes have a partial **prebreeding** moult including body feathers, some scapulars, tertials and central tail feathers.

PHENOLOGY



STATUS EN ARAGON

Summer visitor with rare wintering birds. Breeds in rice fields and waterplaces from the Ebro Valley and Sistema Iberico.



Black-winged Stilt. Adult. Female (22-IV)



Black-winged Stilt. Ageing. Pattern of upperparts: left adult; right juvenile.







Blackwinged Stilt. Ageing. Colour of legs: top adult; bottom juvenile.

http://blascozumeta.com

Put your logo here



Black-winged Stilt. 2nd year. Male ()



Black-winged Stilt. 2nd year. Female ()



Black-winged Stilt. Juvenile (21-VII).




Black-winged Stilt. Young (22-VI)




Blackwinged Stilt. Adult. Head pattern: top male (22-IV); bottom female (22-IV)




Put your logo here





Blackwinged Stilt. Head pattern: top 2nd year male (); middle 2nd year female (); bottom juvenile (21 -VII).







Black-winged Stilt. Nape pattern: top left 2nd year male (); top right 2nd year female (); left juvenile (21-VII).





Black-winged Stilt. Adult. Nape pattern: left male (22-IV); right female (22-IV)



Black-winged Stilt. Adult. Crown pattern: left male (22-IV); right female (22-IV)

Put your logo here





Blackwinged Stilt. Head pattern: top 2nd year male (); middle 2nd year female (); bottom juvenile (21 -VII).



Black-winged Stilt. Crown pattern: top left 2nd year male (); top right 2nd year female (); left juvenile (21-VII).









Blackwinged Stilt. Adult. Breast pattern: top male (22-IV); bottom female (22-IV)




Black-winged Stilt. Adult. Upperpart pattern: left male (22-IV); right female (22-IV)

Put your logo here






Black-winged Stilt. Upperpart pattern: top left 2nd year male (); top right 2nd year female (); left juvenile (21-VII).



Blackwinged Stilt. Legs pattern: top 2nd year male (); middle 2nd year female (); bottom juvenile (21 -VII).







Blackwinged Stilt. Adult. Legs pattern: top male (22-IV); bottom female (22-IV)








Black-winged Stilt. Adult. Tail pattern: left male (22 -IV); right female (22-IV)


http://blascozumeta.com

Put your logo here











Black-winged Stilt. Tail pattern: top left 2nd year male (); top right 2nd year female (); left juvenile (21-VII).



Black-winged Stilt. Pattern of tertials: top left 2nd year male (); top right 2nd year female (); left juvenile (21-VII).



Black-winged Stilt. Adult. Pattern of tertials: left male (22-IV); right female (22-IV)




Black-winged Stilt. Adult. Male: pattern of wing coverts (22-IV).


Put your logo here



Black-winged Stilt. Adult. Female: pattern of wing coverts (22-IV).



Black-winged Stilt. 2nd year. Male: pattern of wing coverts ().



Black-winged Stilt. 2nd year. Female: pattern of wing coverts ().




Black-winged Stilt. Juvenile: pattern of wing coverts (21-VII)



Black-winged Stilt. Adult. Male: pattern of primaries (22-IV)




Black-winged Stilt. Adult. Female: pattern of primaries (22-IV)



Black-winged Stilt. 2nd year. Male: pattern of primaries ()

http://blascozumeta.com

Put your logo here




Black-winged Stilt. 2nd year. Female: pattern of primaries ()



Black-winged Stilt. Juvenile: pattern of primaries (21 -VII)




Black-winged Stilt. Adult. Female: pattern of secondaries (22-IV)





Black-winged Stilt. Adult. Male: pattern of secondaries (22-IV)

Black-winged Stilt. 2nd year. Male: pattern of secondaries ()




Black-winged Stilt. 2nd year. Female: pattern of se-condaries ()


Put your logo here

Black-winged Stilt 162



Black-winged Stilt. Juvenile: pattern of secondaries (21-VII)





Black-winged Stilt. Adult. Male: pattern of wing (22 -IV)







Black-winged Stilt. Adult. Female: pattern of wing (22-IV)

Black-winged Stilt. 2nd year. Female: pattern of wing ()



Black-winged Stilt. Juvenile: pattern of wing (21-VII)