




Woodcock (23-XII).

EURASIAN WOODCOCK (Scolopax rusticola)

IDENTIFICATION

34-36 cm. Grey upperparts; with black and buff streaked; grey underparts, with transversal lines; grey and brown crown, with pale transversal lines; brown wings con pale buff bands; dark tail, with reddish tinge; long bill, darker on tip.



SIMILAR SPECIES

Recalls a Common Snipe (25-28 cm), which is smaller and has longitudinal lines on crown.



SEXING

Plumage of both sexes alike.

AGEING

3 types of age can be recognized:

Juvenile similar to adults but with tips of primary coverts broad buff edged; fresh plumage.

1st year autumn/2nd year spring maintain tips of primary coverts broad buff edged; pale flight feathers; primaries with worn tips.

Adult with narrow pale tips on primary coverts; dark flight feathers; primaries with fresh tips.

CAUTION: wear on primary tips in not evident in early winter.





Wood cock. Ageing. Pattern of tips of primary coverts: top adult; bottom 1st year.

http://blascozumeta.com http://www.aranzadi.eus







Woodcock. Ageing. Pattern of primaries: top adult; bottom 1st year.






Woodcock. Ageing. Pattern of tail feathers (from below): left adult; right 1st year.

MOULT

Complete **postbreeding** moult, usually finished in September. Partial **postjuvenile** moult involving body feathers, all or most of tail, lesser and median coverts and inner secondaries with their coverts; usually finished in October. **Adults** have a **prebreeding** moult including some body feathers, wing coverts and tail feathers.

PHENOLOGY



PICTURE NOT AVAILABLE

Woodcock. Adult ().

PICTURE NOT AVAILABLE

Woodcock. Juvenile ().

PICTURE NOT AVAILABLE



Woodcock. Head pattern: top adult (); middle 1st year (23-XII); bottom juvenile ().

PICTURE NOT AVAILABLE

http://blascozumeta.com http://www.aranzadi.eus



PICTURE NOT AVAILABLE



PICTURE NOT AVAILABLE

Woodcock. Breast pattern: top left adult (); top right 1st year (23-XII); left juvenile ().



PICTURE NOT AVAILABLE



Woodcock. Tail pattern: top left adult (); top right 1st year (23-XII); left juvenile ().

PICTURE NOT AVAILABLE

PICTURE NOT AVAILABLE



Woodcock. Upperpart pattern: top left adult (); top right 1st year (23-XII); left juvenile ().



Woodcock. Adult: pattern of primary coverts (16-XII).



Woodcock. 1st year: pattern of primary coverts (16-XII).

3

http://blascozumeta.com http://www.aranzadi.eus



PICTURE NOT AVAILABLE

Woodcock. Juvenile: pattern of primary coverts ().



Woodcock. Adult: pattern of primaries (16-XII).



Woodcock. 1st year: pattern of primaries (16-XII).

PICTURE NOT AVAILABLE

Woodcock. Juvenile: pattern of primaries ().



Woodcock. Adult: pattern of secondaries (16-XII).



Woodcock. 1st year: pattern of secondaries (16-XII).



PICTURE NOT AVAILABLE

Woodcock. Juvenile: pattern of secondaries ().



Woodcock. Adult: pattern of wing (16-XII).



Woodcock. 1st year: pattern of wing (16-XII).

PICTURE NOT AVAILABLE

Woodcock. Juvenile: pattern of wing ().