



Redshank. Spring. Adult (02-V)

COMMON REDSHANK (Tringa totanus)

IDENTIFICATION


27-28 cm. In **spring** with brownish upperparts barred dark; white rump; white tail with dark bars; wings with a broad white band on the edge; white underparts with dark streaks on head, neck and breast; red bill with dark tip; orange-red legs. In **autumn** with darker colours on upperparts: grey tinge on breast.




Redshank. Pattern of wing, tail and bill.

SIMILAR SPECIES

Recalls a **Spotted Redhsank** in **autumn**, without a white patch on wings and has a longer bill; **Ruff** has a narrower wing band and two white bands on sides of uppertail





SEXING

Plumage of both sexes alike.

AGEING

This species is a scarce **breeder** in Aragon, so only 3 age classes can be recognized:

1st year autumn with fresh plumage; median coverts spotted pale on edge; tertials brown with buff and dark marks; pointed tail feathers; breast slightly streaked; dull reddish base of bill; yellowish legs.

2nd year spring similar to **adult**; this age can be recognized only in birds with some unmoulted median wing coverts and/or tertials; flight feathers moderately worn.

Adult with median coverts with whitish edge and dark subterminal band; tertials grey brown with variable markings: either plain or with dark sepia bars, sometimes with extensive dark markings; rounded tail feathers; underparts with variable amount of dark barring and spotting; reddish legs and base of bill.





Redshank. Ageing. Pattern of bill: top adult; bottom 1st year.







Redshank. Ageing. Pattern of legs: top adult; bottom 1st year.






Redshank. Ageing. Pattern of tertials: left adult in spring; right 1st year.

MOULT

Complete **postbreeding** moult, starting in breeding areas and finished in wintering sites. Partial **postjuvenile** moult involving body feathers, most scapulars, some tail feathers (occasionally all), some tertials and some lesser and median wing coverts; started in breeding areas and fisnished in wintering quarters. Both age classes have a **prebreeding** moult including all or part of body feathers, some scapulars, tertials, wing coverts and sometines some tail feathers.

PHENOLOGY





Redshank. Spring. 2nd year (04-V)





Redshank. Spring. Head pattern: top adult (02-V); bottom 2nd year (04-V)

2







Redshank. Spring. Breast pattern: top adult (02-V); bottom 2nd year (04-V) $\,$





Redshank. Spring. Upperpart pattern: left adult (02-V); right 2nd year (04-V)





Redshank. Spring. Tail pattern: left adult (02-V); right 2nd year (04-V)



Redshank. Spring. Adult: pattern and wear of wing coverts (02-V)



Redshank. Spring. 2nd year: pattern and wear of wing coverts (04-V)







Redshank. Spring. Pattern of tertials: left adult (02-V); right 2nd year (04-V)



Redshank. Spring. Adult: pattern of primaries (02-V)



Redshank. Spring. 2nd year: pattern of primaries (04-V)



Redshank. Spring. Adult: pattern of secondaries (02-V)



Redshank. Spring. 2nd year: pattern of secondaries (04-V)





Redshank. Spring. Adult: pattern of wing (02-V)

http://www.aranzadi.eus







Redshank. Spring. 2nd year: pattern of wing (04-V)



Redshank. Autumn. Adult (07-VII) (Photo: Ottenby Bird Observatory)



Redshank. Autumn. 1st year (04-IX)





Redshank. Autumn. Head pattern: top adult (Photo: Ottenby Bird Observatory); bottom 1st year (04-IX)

PICTURE NOT AVAILABLE

Redshank.
A u t u m n .
Breast pattern: top
adult ();
bottom 1st
year (04-IX)







Redshank. Autumn. Upperpart pattern: left adult (Photo: Ottenby Bird Observatory); right 1st year (04-IX)



PICTURE NOT AVAILABLE



Redshank. Autumn. Legs pattern: top adult (); bottom 1st year (04-IX)





Redshank. Autumn. Tail pattern: left adult (Photo: Ottenby Bird Observatory); right 1st year (04-IX)



Redshank. Autumn. Adult: pattern of wing coverts (Photo: Ottenby Bird Observatory)



Redshank. Autumn. 1st year: pattern of wing coverts (04-IX)





Redshank. Autumn. Pattern of tertials: left adult (Photo: Ottenby Bird Observatory); right 1st year (04-





Redshank. Autumn. Adult: pattern of primaries (Photo: Ottenby Bird Observatory)



Redshank. Autumn. 1st year: pattern of primaries (04 -IX)



Redshank. Autumn. Adult: pattern of secondaries (Photo: Ottenby Bird Observatory)



Redshank. Autumn. 1st year: pattern of secondaries (04-IX)



PICTURE NOT AVAILABLE

Redshank. Autumn. Adult: pattern of wing (Photo: Ottenby Bird Observatory)





Redshank. Autumn. 1st year: pattern of wing (04-IX)