

Stock Dove. Spring. Male (28-IV)

STOCK DOVE (Columba oenas)

IDENTIFICATION

30-33 cm. Adult with bluish grey plumage, lacking white on wing or rump; wing with two black bands; green and purple patch on neck; grey tail with two terminal bands; reddish legs; red bill with yellow tip. Juvenile without green and purple patch on neck.







Stock Dove. Pattern of rump, underwing coverts and wing.

SIMILAR SPECIES

Recalls a **Feral Pigeon/Rock Dove** which have white rump and underwing coverts. **Wood Pigeon** has a bigger size and has a white transverse band on wings.





SEXING

Male with bluish grey mantle, tertials and scapulars. Female with brownish grey mantle, tertials and scapulars (CAUTION: with worn plumage both sexes are difficult to separe). Juveniles cannot be sexed using plumage pattern





Stock Dove.
Sexing. Pattern of iridescent patch on neck: top male; bottom female.










Stock Dove. Ageing. Pattern of head and neck: top adult; bottom juvenile.











Stock Dove. Ageing. Pattern of legs: adult; top bottom juveni-

Stock Dove. Sexing. Pattern of scapulars: left male; right female.

AGEING

4 age groups can be recognized:

Juvenile without iridescence on neck; mantle feathers, wing coverts and tertials brownish; light brownish primary coverts; pinkish legs.

1st year autumn recognized till moult is completed by unmoulted flight and tail feathers only slightly worn.

1st year autumn/2nd year spring only birds with arrested moult, which have pale brown juvenile outer primary coverts.

Adult with iridescence on neck; mantle feathers, wing coverts and tertials bluish-grey; greyish black primary coverts; reddish legs. In autumn recognizable till moult is completed by very worn unmoulted flight and tail feathers. In spring, birds with arrested moult can be recognized by their greyish black primary coverts.





Stock Dove. Ageing. Pattern of primary coverts: adult; top bottom juvenile.





Stock Dove. Autumn. Ageing. Wear of unmoulted tail feathers: left adult; right 1st year.





Stock Dove. Autumn. Ageing. Wear of unmoulted secondaries: top adult; bottom 1st year.





Stock Dove. Autumn. Ageing. Wear of unmoulted primaries: top adult; bottom 1st year.





Stock Dove. Autumn. Ageing. Wear of wing feathers: top adult; bottom 1st year.

MOULT


Complete **postbreeding** and **postjuvenile** moults, usually finished in December (later in late brood **juveniles**). Both age classes, more frequent in **juvenile** birds, may arrested wing moult and have unmoulted flight feathers.

PHENOLOGY

| I | II | III | IV | V | VI | VII | VIII | IX | X | XI | XII |
|---|----|-----|----|---|----|-----|------|----|---|----|-----|
| | | | | | | | | | | | |

STATUS IN ARAGON

Resident. Widely distributed throughout the Region, absent from the mountainous zones.



Stock Dove. Spring. 2nd year. Female ().



Stock Dove. Spring. Female (17-V).



Stock Dove. Spring. Juvenile (26-VII)




Stock Dove. Spring. 2nd year. Male ().





Stock Dove. Spring. Adult. Head pattern: top male (02-VI); bottom female (17-V).





Stock Dove. Spring. Head pattern: top 2nd year male (08-V); middle 2nd year female (); bottom juvenile (26-VII).





Stock Dove. Spring. Adult. Uppepart pattern: left male (02-VI); right female (17-V).








Stock Dove. Spring. Adult. Breast pattern: left male (02-VI); right female (17-V).











Stock Dove. Spring. Breast pattern: top left 2nd year male (08-V); top right 2nd year female (); left juvenile (06-VI).



Stock Dove. Spring. Upperpart pattern: top left 2nd year male (08-V); top right 2nd year female (); left juvenile (06-VI).





Stock Dove. Spring. Legs pattern: top adult (17-V); bottom juvenile (06 -VI).






Stock Dove. Spring. Adult. Pattern of tertials: left male (02-VI); right female (17-V).





Stock Dove. Spring. Adult. Tail pattern: left male (02-VI); right female (17-V).








Stock Dove. Spring. Tail pattern: top left 2nd year male (08-V); top right 2nd year female (); left juvenile (26-VII).








Stock Dove. Spring. Pattern of scapulars: top left 2nd year male (08-V); top right 2nd year female (); left juvenile (06-VI).



Stock Dove. Spring. Adult. Male: pattern of primary coverts (02-VI).



Stock Dove. Spring. 2nd year. Female: pattern of primary coverts ().



Stock Dove. Spring. Adult. Female: pattern of primary coverts (17-V).



Stock Dove. Spring. Juvenile: pattern of primary coverts (26-VII)



Stock Dove. Spring. 2nd year. Male: pattern of primary coverts (08-V).




Stock Dove. Spring. Adult. Male: pattern of primaries (02-VI).



Stock Dove. Spring. Adult. Female: pattern of primaries (17-V).



Stock Dove. Spring. 2nd year. Male: pattern of primaries (08-V).



Stock Dove. Spring. 2nd year. Female: pattern of primaries ().



Stock Dove. Spring. Juvenile: pattern of primaries (26-VII).




Stock Dove. Spring. Adult. Male: pattern of secondaries (02-VI).



Stock Dove. Spring. Adult. Female: pattern of secondaries (17-V).



Stock Dove. Spring. 2nd year. Male: pattern of secondaries (no. 1: juvenile feathers) (08-V)



Stock Dove. Spring. 2nd year. Female: pattern of secondaries ()



Stock Dove. Spring. Juvenile: pattern of secondaries (06-VI)




Stock Dove. Spring. Adult. Male: pattern of wing (28-IV)



Stock Dove. Spring. Adult. Female: pattern of wing (17-V)



Stock Dove. Spring. 2nd year. Male: pattern of wing (no. 1: juvenile feathers) (08-V)



Stock Dove. Spring. 2nd year. Female: pattern of wing ()



Stock Dove. Spring. Juvenile: pattern of wing (28-IV)



Stock Dove. Autumn. Adult (28-X)



Stock Dove. Autumn. 1st year (28-X)



Stock Do-Au-

X); bottom

pat-

top adult (28-

year

ve. tumn. Head

tern:

1st

(28-X).





Stock Dove. Autumn. Breast pattern: left adult (28-X); right 1st year (28-X).





Stock Dove. Autumn. Upperpart pattern: left (28-X); right 1st year (28-X).





Stock Do-Auve. tumn. Colour of legs: top adult (28-X); bottom 1st year (28-X).





Stock Dove. Autumn. Tail pattern: left adult (28-X); right 1st year (28-X).



Stock Dove. Autumn. 1st year: pattern of primary coverts (28-X).





Stock Dove. Autumn. Pattern of tertials: left adult (28-X); right 1st year (28-X).





Stock Dove. Autumn. Adult: pattern of primaries (28 -X).



Stock Dove. Autumn. Adult: pattern of primary coverts (28-X).





Stock Dove. Autumn. 1st year: pattern of primaries (28-X).





Stock Dove. Autumn. Adult: pattern of secondaries (28-X).





Stock Dove. Autumn. 1st year: pattern of secondaries (28-X).





Stock Dove. Autumn. Adult: pattern of wing (28-X).





Stock Dove. Autumn. 1st year: pattern of wing (28-X)