

Hoopoe. Adult. Male (27-V).

EURASIAN HOOPOE (*Upupa epops*)

IDENTIFICATION

25-28 cm. Brownish upperparts, with transversal black and white lines; pink-vinaceous breast; white underparts; rufous head with a pink crest with black and white tips; wings and tail black with white bands; long and curved bill.

SIMILAR SPECIES

This species is unmistakable.

AGEING

3 age groups can be recognized:

Juvenile resembles a **female**, but with fresh flight feathers and without moult limits; outermost tail feather with rounded tip.

2nd year with unmoulted flight feathers very worn; if tertials and inner secondaries changed then with moult limit between these darker and fresh feathes and the worn and bleached rest of retained juvenile flight feathers; if tail is unmoulted, then with very worn feathers.

Adult without moult limit on wing feathers, which will be fresh; outermost tail feather with squared tip.





Hoopoe. Ageing. Pattern of tip of the outermost tail feather: left adult (27-IV); right juvenile (04-VIII).

SEXING

In adult birds, male with chin and breast deep vinaceous pink; mantle lacks pinkish tinge. Female with chin and breast cinnamon rufous; mantle with pale pinkish tinge. Juveniles can be sexed in some extreme birds: males with head and breast more intense pink than females. CAUTION: with worn plumage, colour differences are very difficult to see and sexing is possible only by comparing within pair.

MOULT

Complete **postbreeding** moult, changing in breeding areas body feathers and sometimes some flight and tail feathers between July-August and finished in wintering sites. Partial **postjuvenile** moult involving only body feathers, with some birds changing all or most tail feathers, tertials or inner secondaries; usually moulted in wintering sites. Mainly **adults**, and some **juveniles**, have a **prebreeding** moult including only head and neck feathers.

PHENOLOGY



STATUS IN ARAGON

Summer visitor with some scarce wintering birds. Breeds widely distributed throughout the Region, absent only from Northern areas.



Hoopoe. Adult. Female (17-VI).



Hoopoe. 2nd year. Male (18-IV).



Hoopoe. 2nd year. Female (07-V).



Hoopoe. Juvenile. Male (04-VIII).



Hoopoe. Juvenile. Female (26-VII).



Hoopoe. Adult. Head pattern: top male (27-IV); bottom female (17-VI).

Page 2





Hoopoe. 2nd year. Head pattern: top male (18-IV); bottom female (07-V).





Hoopoe. 2nd year. Breast pattern: left male (18-IV); right female (07-V).





Hoopoe. Juvenile. Breast pattern: left male (04-VIII); right female (26-VII).



Hoopoe. Juvenile. Head pattern: top male (04-VIII); bottom female (26-VII).





Hoopoe. Adult. Nape pattern: left male (27-IV); right female (17-VI).





Hoopoe. Adult. Breast pattern: left male (27-IV); right female (17-VI).





Hoopoe. 2nd year. Nape pattern: left male (18-IV); right female (07-V).





Hoopoe. 2nd year. Upperpart pattern: left male (18-IV); right female (07-V).





Hoopoe. Juvenile. Nape pattern: left male (04-VIII); right female (26-VII).





Hoopoe. Juvenile. Upperpart pattern: left male (04-VIII); right female (26-VII).





Hoopoe. Adult. Upperpart pattern: left male (27-IV); right female (17-VI).





Page 4

Hoopoe. Adult. Tail pattern: left male (27-IV); right female (17-VI).



Hoopoe. 2nd year. Tail pattern: left male (27-V); right female (07-V).



Hoopoe. Adult. female: pattern of primaries (17-VI).





Hoopoe. Juvenile. Tail pattern: left male (04-VIII); right female (26-VII).



Hoopoe. 2nd year. Male: pattern of primaries (18-IV).



Hoopoe. Adult. Male: pattern of primaries (27-IV).



Hoopoe. 2nd year. Female: pattern of primaries (07-V).



Hoopoe. Juvenile. Male: pattern of primaries (04-VIII).



Hoopoe. Adult. Female: pattern of secondaries (17-VI).



Hoopoe. Juvenile. Female: pattern of primaries (26-VII).




Hoopoe. 2nd year. Male: pattern of secondaries (18-IV).



Hoopoe. Adult. Male: pattern of secondaries (27-IV).



Hoopoe. 2nd year. Female: pattern of secondaries (07-V).



Hoopoe. Juvenile. Male: pattern of secondaries (04-VIII).



Hoopoe. Juvenile. Female: pattern of secondaries (26 -VII).



Hoopoe. Adult. Male: pattern of wing (27-IV).



Hoopoe. Adult. Female: pattern of wing (17-VI).



Hoopoe. 2nd year. Male: pattern of wing (18-IV).



Hoopoe. 2nd year. Female: pattern of wing (07-V).



Hoopoe. Juvenile. Male: pattern of wing (04-VIII)

Put your logo here



Hoopoe. Juvenile. Female: pattern of wing (26-VII)