

Yellow Wagtail. Spring. Adult. Male (22-IV).

WESTERN YELLOW WAGTAIL (Motacilla flava)

IDENTIFICATION

16-17 cm. Yellowish underparts; brown-olive upperparts; long tail, brown with white edges; dark legs. In subspecies *iberiae*, male in breeding plumage with grey head and white supercilium. Females and juveniles are duller.



Yellow Wagtail. Pattern of upperparts and legs.



SIMILAR SPECIES

Recalls a **Grey Wagtail**, but this species has grey upperparts and pale legs.



SEXING

In breeding plumage, male with grey-bluish crown; white supercilium; underparts deep yellow. Female with grey-brown crown; dull white supercilium; underparts dull yellow with some dark spots on breast. After postbreeding moult, male with bright yellow underparts and grey bluish crown; female with throat and breast whitish buff; dull yellow underparts. After postjuvenile moult, 1st year male is very similar to adult female so it is necessary first a correct ageing; 1st year males have underparts more yellow than 1st year females, which have whitish underparts. CAUTION: last hatched males moult later and resemble females.





Yellow Wagtail. Spring. Sexing. Pattern of head: top male; bottom female.

AGEING

4 types of age can be recognized:

Juvenile with fresh plumage; dull colours; dark narrow collar on upper breast.

1st year autumn with moult limit between moulted fresh greater covers and tertials and worn whitish **juvenile** retained; if all greater coverts are unmoulted limit is between median and lesser coverts; if all greater coverts have been moulted limit is between tertials; tail feathers worn; sometimes without yellow.

2nd year spring similar to **adult**, so ageing is only possible in birds which retaining **juvenile** greater coverts so little worn that they show the white edges on the tip.

Adult in autumn without moult limits; with fresh flight feathers and greater coverts with olive yellowish edges and tips; in spring with greater coverts with diffuse edges and not white. CAUTION: after prebreeding moult ageing is not always possible.





Yellow Wagtail. Spring. Ageing. Pattern of head and breast: left adult female; right juvenile.





Yellow Wagtail. Autumn. Ageing and sexing. Adult. Pattern of head: top male; bottom female.





Yellow Wagtail. Autumn. Ageing and sexing. 1st year. Pattern of head: top male; bottom female.





Yellow Wagtail. Autumn. Ageing and sexing. Adult. Pattern of undertail coverts: left male; right female.






Yellow Wagtail. Autumn. Ageing and sexing. 1st year. Pattern of undertail coverts: left male; right female.

MOULT

Complete postbreeding moult, usually finished late September; some birds can suspend the moult and finish it in their wintering quarters. Partial postjuvenile moult including body feathers and lesser and median coverts; some birds can change some or all greater coverts and some or all tertials; usually finished early September. Between February and April, both types of age have a prebreeding moult acquiring breeding plumage, changing all or some inner greater coverts, tertials and tail feathers.



Yellow Wagtail. Extent of postjuvenile moult.

PHENOLOGY



STATUS IN ARAGON

Summer visitor. Very common on passage. Breeds widely distributed throughout the Region.



Yellow Wagtail. Spring. Adult. Female (26-IV)



Yellow Wagtail. Spring. 2nd year. Male (02-IV)



Yellow Wagtail. Spring. 2nd year. Female (02-IV)



Yellow Wagtail. Juvenile (30-VII)



Yellow Wagtail. Spring. Adult. Head pattern: top male (26-IV); bottom female (26-IV).









Yellow Wagtail. Spring. Breast pattern: top left 2nd year male (02-IV); top right 2nd year female (02-IV); left juvenile (30-VII).





Yellow Wagtail. Spring. Head pattern: top 2nd year male (02-IV); middle 2nd year female (02-IV); botom juvenile (30-VII).











Yellow Wagtail. Spring. Adult. Breast pattern: left male (26-IV); right female (26-IV).

Yellow Wagtail. Spring. Adult. Upperparts pattern: left male (26-IV); right female (26-IV).







Yellow Wagtail. Spring. Upperparts pattern: top left 2nd year male (02-IV); top right 2nd year female (02-IV); left juvenile (30-VII).








Yellow Wagtail. Spring. Underparts pattern: top left 2nd year male (02-IV); top right 2nd year female (02-IV); left juvenile (30-VII).





Yellow Wagtail. Spring. Adult. Underparts pattern: left male (26-IV); right female (26-IV).







Yellow Wagtail. Spring. Pattern of undertail coverts: top left 2nd year male (02-IV); top right 2nd year female (02-IV); left juvenile (30-VII).





Yellow Wagtail. Spring. Adult. Pattern of undertail coverts: left male (26-IV); right female (26-IV).



Yellow Wagtail. Spring. Adult. Male: pattern of wing (02-IV).



Yellow Wagtail. Spring. Adult. Female: pattern of wing (22-IV).



Yellow Wagtail. Autumn. Adult. Male (01-IX).



Yellow Wagtail. Spring. 2nd year. Male: pattern of wing (22-IV).



Yellow Wagtail. Autumn. Adult. Female (14-IX).



Yellow Wagtail. Spring. 2nd year. Female: pattern of wing (02-IV).



Yellow Wagtail. Spring. Juvenile: pattern of wing (30-VII).



Yellow Wagtail. Autumn. 1st year. Male (14-XI).



Yellow Wagtail. Autumn. 1st year. Female (14-IX).



Yellow Wagtail. Autumn. Adult. Head pattern: top male (14-IX); bottom female (01-IX).







Yellow Wagtail. Autumn. 1st year. Head pattern: top male (01-IX); bottom female (27-VIII).





Yellow Wagtail. Autumn. Adult. Breast pattern: left male (01-IX); right female (14-IX).





Yellow Wagtail. Autumn. 1st year. Breast pattern: left male (01-IX); right female (27-VIII).





Yellow Wagtail. Autumn. Adult. Pattern of undertail coverts: left male (01-IX); right female (27-VIII).



Yellow Wagtail. Autumn. 1st year. Pattern of undertail coverts: left male (01-IX); right female (01-IX).





Yellow Wagtail. Autumn. 1st year. Upperparts pattern: left male (27-VIII); right female (27-VIII).





Yellow Wagtail. Autumn. Adult. Upperparts pattern: left male (01-IX); right female (27-VIII).





Yellow Wagtail. Autumn. Adult. Underparts pattern: left male (01-IX); right female (27-VIII).





Yellow Wagtail. Autumn. 1st year. Underparts pattern: left male (14-IX); right female (27-VIII).



Yellow Wagtail. Autumn. Adult. Male: pattern of wing (01-IX).



Yellow Wagtail. Autumn. Adult. Female: pattern of wing (27-VIII).



Yellow Wagtail. Autumn. 1st year. Male: pattern of wing (27-VIII).



Yellow Wagtail. Autumn. 1st year. Female: pattern of wing (27-VIII).