



Dipper. Adult (8-I).

WHITE-THROATED DIPPER (Cinclus cinclus)

IDENTIFICATION

16-19 cm. Adult with dark grey upperparts; redbrown head and belly; white breast and throat. Juvenile dull grey.



Dipper. Adult. Pattern of

head and upperparts.

SIMILAR SPECIES

This species is unmistakable.

SEXING

Plumage of both sexes alike. Size is helpful with most of the specimens: male with wing longer than 94 mm; female with wing shorter than 89 mm.

AGEING

3 types of age can be recognized:

Juvenile dull grey, with grey wavy bars on underparts; primary coverts narrow and pointed; grey-olive

1st year autumn/2nd year spring with moult limit between the moulted wing coverts and the juvenile retained with white tips.

Adult with all wing coverts dark, without white tips; primary coverts broad and rounded; brown-reddish iris.





Dipper. Ageing. Pattern of head and iris colour: top adult; bottom juvenile.





Dipper. Ageing. Pattern of breast: left adult; right juvenile.





Dipper. Ageing. Moult limit on wing coverts: top adult; bottom 1st year.


1

http://blascozumeta.com http://www.aranzadi.eus




MOULT

Complete **postbreeding** moult, often finished in September. Partial **postjuvenile** moult including body feathers and lesser and median wing coverts; some birds moult a variable number of inner greater coverts; usually finished in October.



D i p p e r . Extent of postjuvenile moult.

PHENOLOGY





Dipper. 1st year (27-XII).



Dipper. Juvenile (12-VII).







Dipper. Head pattern and iris colour: top adult (27-XII); middle 1st year (27-XII); bottom juvenile (12-VII).







Dipper. Breast pattern: top left adult (27-XII); top right 1st year (27-XII); left juvenile (12-VII).

2

http://blascozumeta.com http://www.aranzadi.eus









Dipper. Upperpart pattern: top left adult (27-XII); top right 1st year (27-XII); left juvenile (12-VII).







Dipper. Tail pattern: top left adult (27-XII); top right 1st year (27-XII); left juvenile (12-VII).



Dipper. Adult: pattern of primary coverts (08-I).



Dipper. 1st year: pattern of primary coverts (27-XII).



Dipper. Juvenile: pattern of primary coverts (12-VII).

http://blascozumeta.com http://www.aranzadi.eus 3





Dipper. Adult: pattern of wing (27-XII).



Dipper. 1st year: pattern of wing (27-XII).



Dipper. Juvenile: pattern of wing (12-VII).