

324 Dunnock

[Put your logo here](#)




Dunnock. Adult (06-I).


DUNNOCK (*Prunella modularis*)

IDENTIFICATION

13-15 cm. Grey head, neck and breast; brown and streaked upperparts; pale underparts.




Dunnock. Pattern of head, breast and upperparts.



SIMILAR SPECIES

Recalls an **Alpine Accentor**, with barred throat and chestnut flanks.



Alpine Accentor

SEXING

Male usually with head and breast more lead-grey than **female**, but it is not an useful characteristic for sexing due to overlap.

AGEING

3 types of age can be recognized:


Juvenile with fresh plumage; head and breast streaked; greyish iris.

1st year autumn/2nd year spring with unmoulted **juvenile** greater coverts with pale marks on tips of both webs and distinct black central tip; if there are moulted inner greater coverts, then with indistinct pale marks shading into black central tip; unmoulted alula feathers dull; bill with base of the lower mandible paler; pointed tail feathers; dull brown iris (**CAUTION:** some **juvenile** birds have reddish tinge on iris in early October).



Adult without moult limit on wings; outer greater coverts with pale marks on the tips, while the inner ones do not have pale marks or are small; alula feathers dark; reddish iris; dark bill; rounded tail feathers.

324 Dunnock

Put your logo here




Dunnock. Ageing. Pattern of head: top adult; bottom juvenile.



Dunnock. Ageing. Pattern of tips of tail feathers: left adult; right 1st year.




Dunnock. Ageing. Pattern of breast: left adult; right juvenile.



Dunnock. Ageing. Colour of iris and bill: top adult; middle 1st year; bottom juvenile.

MOULT

Complete **postbreeding** moult, usually finished early October. Partial **postjuvenile** moult including body feathers and lesser and median wing coverts, but some birds may extending their moult to a variable number of greater coverts, tertials and tail feathers; usually finished late September.



Dunnock. Extent of postjuvenile moult.

PHENOLOGY

I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XII

STATUS IN ARAGON

Resident, breeds in mountainous areas with presence in lower areas in winter and contribution of wintering European birds.

324 Dunnock

Put your logo here



Dunnock. 1st year (02-XII).




Dunnock. Breast pattern: top left adult (01-XI); top right 1st year (02-XII); left juvenile (18-VII).



Dunnock. Juvenile (18-VII)



Dunnock. Tail pattern: top left adult (06-I); top right 1st year with fresh tail (02-XII); left 1st year with worn tail (02-XII); bottom juvenile (18-VII).




Dunnock. Head pattern and iris colour: top adult (06-I); middle 1st year (02-XII); bottom juvenile (18-VII).



324 Dunnock


Put your logo here



Dunnock. Legs pattern: left adult (06-XII); right juvenile (18-VII).



Dunnock. Adult: pattern of alula (06-XII).



Dunnock. 1st year: pattern of alula (02-XII).



Dunnock. Juvenile: pattern of alula (18-VII).



Dunnock. Upperpart pattern: top left adult (06-I); top right 1st year (02-XII); left juvenile (18-VII).



Dunnock. Adult: pattern of wing (06-XII).

324 Dunnock

Put your logo here



Dunnock. 1st year: pattern of wing (02-XII).



Dunnock. Juvenile: pattern of wing (18-VII).