

325 Alpine Accentor

[Put your logo here](#)



Alpine Accentor. Adult (22-II).


ALPINE ACCENTOR (*Prunella collaris*)

IDENTIFICATION

16-18 cm. Grey upperparts with brown spots; greyish underparts with chestnut flanks; white throat with brown bars; wings with dark coverts tipped white; brown tail with white tips.




Alpine Accentor.
Pattern of head,
breast and upper-
parts.



SIMILAR SPECIES

Recalls a **Dunnock**, without barred throat and chestnut flanks.



Dunnock



SEXING

Plumage of both sexes alike. Size can be an useful characteristic in extreme birds: **male** with wing longer than 109 mm; **female** with wing shorter than 96 mm.

AGEING

3 types of age can be recognized:

Juvenile with very fresh plumage; buff underparts with bad defined streaked; buff flanks with dark streaked; iris brown-greyish.

1st year autumn/2nd year spring with moult limit between median and greater coverts or within greater coverts; unmoulted **juvenile** greater coverts with a distal brown-white mark broad and rounded; with a narrow edge between pale mark and brown-white tip; flight and tail feathers eroded.

Adult without moult limits on the wing; greater coverts with pure white marks on tips, contrasting with the black bases of feathers; unstreaked grey underparts; with chestnut flanks; flight and tail feathers fresh; iris brown-reddish.

325 Alpine Accentor

Put your logo here



Alpine Accentor. Ageing. Pattern of head and colour of iris: top adult; bottom juvenile




Alpine Accentor. Ageing. Pattern of breast: left adult; right juvenile



Alpine Accentor. Ageing. Pattern of greater coverts: top adult; bottom 2nd year

MOULT

Complete **postbreeding** moult, usually finished in November. Partial **postjuvenile** moult including body feathers and lesser and median wing coverts, but some birds may extend their moult to a variable number of inner greater coverts and tertials; usually finished in October.



Alpine Accentor. Extent of postjuvenile moult.

■ Todos los ejemplares
■ Algunos ejemplares

PHENOLOGY

I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XII

STATUS IN ARAGON

Resident, breeds in mountainous areas with wide presence in the Pyrenees and quite scarce in Moncayo.



Alpine Accentor. 2nd year (06-II).

325 Alpine Accentor

Put your logo here




Alpine Accentor. Juvenile (05-VII).



Alpine Accentor. Breast pattern: top left adult (22-II); top right 2nd year (06-II); left juvenile (05-VII).




Alpine Accentor. Head pattern and iris colour: top adult (22-II); middle 2nd year (06-II); bottom juvenile (05-VII).



Alpine Accentor. Pattern of upperparts: top left adult (22-II); top right 2nd year (06-II); left juvenile (05-VII).

325 Alpine Accentor

Put your logo here




Alpine Accentor. Tail pattern: top left adult (22-II); top right 2nd year (06-II); left juvenile (05-VII).



Alpine Accentor. Juvenile: pattern of flank (05-VII)



Alpine Accentor. Adult: pattern of greater coverts (22-II).



Alpine Accentor. Adult: pattern of flank (22-II)



Alpine Accentor. 2nd year: pattern of greater coverts (06-II).



Alpine Accentor. 2nd year: pattern of flank (06-II)

325 Alpine Accentor

[Put your logo here](#)



Alpine Accentor. Juvenile: pattern of greater coverts (05-VII).



Alpine Accentor. Juvenile: pattern of primary coverts (05-VII).



Alpine Accentor. Adult: pattern of primary coverts (22-II).



Alpine Accentor. Adult: pattern of wing (22-II).



Alpine Accentor. 2nd year: pattern of primary coverts (06-II).



Alpine Accentor. 2nd year: pattern of wing (06-II).



Alpine Accentor. Juvenile: pattern of wing (05-VII).