

Moustached Warbler. Spring (10-III).

MOUSTACHED

WARBLER

(Acrocephalus melanopogon)

IDENTIFICATION

11-13 cm. Brown reddish upperparts, streaked black; unstreaked nape and rump; distinct white supercilium; dark crown; white underparts, with buff throat; brown tail with reddish edges.



SIMILAR SPECIES

Recalls a **Reed Warbler**, without pale supercilium. Similar to **Aquatic Warbler** with a pale band along crown. **Sedge Warbler** has dark streaked crown, narrower pale supercilium and buff throat.



Reed Warbler





SEXING

Plumage of both sexes alike.

AGEING

2 types of age can be recognized:

Juvenile with fresh feathers (check flight feathers and tail); iris brown-olive; yellow inside of upper mandible; 10th primary with broad tip; with two black tongue spots.

Adult in autumn with worn plumage; iris brown-reddish; orange inside of upper mandible; 10th primary with narrow tip; without tongue spots.

After **postbreeding/postjuvenile** moults, ageing is not possible using plumage pattern.





Moustached Warbler. Summer. Ageing. Wear of tertials: left adult; right juvenile.





Moustached Warbler. Summer. Ageing. Colour of iris: top adult; bottom juvenile.

MOULT

Complete **postbreeding** and **postjuvenile** moults, usually finished in October.

PHENOLOGY



STATUS IN ARAGON

Resident. Breeding only in Laguna de Sariñena.





Moustached Warbler. Summer. Ageing. Colour inside of upper mandible: left adult; right juvenile.



Moustached Warbler. Summer. Adult (02-VII).





Moustached Warbler. Summer. Ageing. Pattern of tip of the 10th primary: top adult; bottom juvenile.



Moustached Warbler. Summer. Juvenile (02-VI).





Moustached Warbler. Head pattern and iris colour: top in spring (10-III); middle adult in summer (02-VI); bottom juvenile (02-VI).





Moustached Warbler. Crown pattern: top left in spring (10-III); top right adult in summer (02-VI); left juvenile (02-VI).









Moustached Warbler. Breast pattern: top left in spring (10-III); top right adult in summer (02-VI); left juvenile (02-VI).







Moustached Warbler. Upperpart pattern: top left in spring (10-III); top right adult in summer (02-VI); left juvenile (02-VI).







Moustached Warbler. Tail pattern: top left in spring (10-III); top right adult in summer (02-VI); left juvenile (02-VI).



Moustached Warbler. Spring: wear of primaries and pattern of 10th primary (10-III).



Moustached Warbler. Summer. Adult: wear of primaries and pattern of 10th primary (02-VI).



Moustached Warbler. Summer. Juvenile: wear of primaries and pattern of 10th primary (02-VI).



Moustached Warbler. Spring: pattern of wing (10-III).



Moustached Warbler. Summer. Adult: pattern of wing (02-VI).



Moustached Warbler. Summer. Juvenile: pattern of wing (02-VI).