

Moustached Warbler. Spring (10-III).

**MOUSTACHED WARBLER** (Acrocephalus melanopogon)

#### **IDENTIFICATION**

11-13 cm. Brown reddish upperparts, streaked black; unstreaked nape and rump; distinct white supercilium; dark crown; white underparts, with buff throat; brown tail with reddish edges.



Moustached Warbler. Pattern of head and crown.

#### SIMILAR SPECIES

Recalls a Reed Warbler, without pale supercilium. Similar to Aquatic Warbler with a pale band along crown. Sedge Warbler has dark streaked crown, narrower pale supercilium and buff throat.



Reed Warbler





Aquatic Warbler

### SEXING

Plumage of both sexes alike.

#### AGEING

2 types of age can be recognized:

Juvenile with fresh feathers (check flight feathers and tail); iris brown-olive; yellow inside of upper mandible; 10th primary with broad tip; with two black tongue spots.

Adult in autumn with worn plumage; iris brownreddish; orange inside of upper mandible; 10th primary with narrow tip; without tongue spots.

After postbreeding/postjuvenile moults, ageing is not possible using plumage pattern.



Moustached Warbler. Summer. Ageing. Wear of tertials: left adult; right juvenile.

http://blascozumeta.com

Sedge Warbler

aranzadi







Moustached Warbler. Summer. Ageing. Colour of iris: top adult; bottom juvenile.

### MOULT

Complete **postbreeding** and **postjuvenile** moults, usually finished in October.

#### PHENOLOGY



Summer visitor
Wintering
Resident
On passage



Moustached Warbler. Summer. Ageing. Colour inside of upper mandible: left adult; right juvenile.



Moustached Warbler. Summer. Adult (02-VII).





Moustached Warbler. Summer. Ageing. Pattern of tip of the 10th primary: top adult; bottom juvenile.



Moustached Warbler. Summer. Juvenile (02-VI).

http://blascozumeta.com







Moustached Warbler. Head pattern and iris colour: top in spring (10-III); middle adult in summer (02-VI); bottom juvenile (02-VI).







Moustached Warbler. Crown pattern: top left in spring (10-III); top right adult in summer (02-VI); left juvenile (02 -VI).







Moustached Warbler. Breast pattern: top left in spring (10-III); top right adult in summer (02-VI); left juvenile (02 -VI).







Moustached Warbler. Upperpart pattern: top left in spring (10-III); top right adult in summer (02-VI); left juvenile (02-VI).









Moustached Warbler. Summer. Adult: wear of primaries and pattern of 10th primary (02-VI).



Moustached Warbler. Tail pattern: top left in spring (10-III); top right adult in summer (02-VI); left juvenile (02-VI).





Moustached Warbler. Spring: wear of primaries and pattern of 10th primary (10-III).

Moustached Warbler. Summer. Juvenile: wear of primaries and pattern of 10th primary (02-VI).



Moustached Warbler. Spring: pattern of wing (10-III).



Moustached Warbler. Summer. Adult: pattern of wing (02-VI).



Moustached Warbler. Summer. Juvenile: pattern of wing (02-VI).

vientzia elkartea