

378a Yellow-browed Warbler

[Put your logo here](#)



Yellow-browed Warbler. Autumn. Adult (11-X)
(Photo: Pere Josa)

YELLOW-BROWED WARBLER (*Phylloscopus inornatus*)

IDENTIFICATION

9-10,5 cm. Greyish-green upperparts; yellow supercilium; off-white underparts with yellowish tinge; white tipped tertials and secondaries; wing with two yellowish bars.



Yellow-browed Warbler. Pattern of tertials and wing-bars

SIMILAR SPECIES

In Aragon this species is unmistakable due to pattern of tertials and wing-bars

SEXING

Plumage of both sexes alike.

AGEING


This is a divagant species in Aragon found mainly in autumn, so only 2 age groups can be recognized:

1st year autumn with greenish upperparts; with contrast between moulted median coverts, with dark centre and unmoulted greater coverts with brown centre; if median coverts unmoulted then with brown tinge; tail feathers pointed and with brown tinge; secondaries with rounded tips and brownish tinge.

Adult with grey-greenish upperparts; both median and greater coverts with dark centre without contrast; tail feathers with broad tips and blackish tinge; secondaries with square tips and blackish tinge.




Yellow-browed Warbler. Autumn. Ageing. Upperpart pattern (Pgoto: Toni Polo Aparisi): left adult; right 1st year



Yellow-browed Warbler. Autumn. Ageing. Tail pattern: left adult (Photo: Pere Josa); right 1st year.

378a Yellow-browed Warbler

Put your logo here



Yellow-browed Warbler. Autumn. Ageing. Pattern of wing coverts: top adult (Photo: Pere Josa); bottom 1st year.



Yellow-browed Warbler. Autumn. Ageing. Pattern of secondaries: left adult (Photo: Pere Josa); right 1st year.

MOULT

Complete **postbreeding** moult, usually finished in August. Partial **postjuvenile** moult involving body feathers and sometimes lesser and median coverts and some tail feathers.

PHENOLOGY

| I | II | III | IV | V | VI | VII | VIII | IX | X | XI | XII |
|---|----|-----|----|---|----|-----|------|----|---|----|-----|
| | | | | | | | | | | | |

STATUS IN ARAGON



An irregular and scarce visitor that can be found anywhere.



Yellow-browed Warbler. Autumn. 1st year (28-X).



Yellow-browed Warbler. Autumn. Head pattern: top adult (11-X) (Photo: Pere Josa); bottom 1st year (28-X).



Yellow-browed Warbler. Autumn. Breast pattern: left adult (18-I) (Photo: Begoña Varcárcel); right 1st year (28-X).

378a Yellow-browed Warbler

Put your logo here



Yellow-browed Warbler. Autumn. 1st year: pattern of wing (28-X).

REFERENCE

Zucca, M., 2017. Évolution récente du statut du Pouillot à grands sourcils *Phylloscopus inornatus* en France. *Ornithos*, 24-4: 201-223

Yellow-browed Warbler. Autumn. Upperpart pattern: left adult (08-XI) (Photo: Toni Polo Aparisi); right 1st year (28-X).



Yellow-browed Warbler. Autumn. Tail pattern: left adult (11-X) (Photo: Pere Josa); right 1st year (28-X).



Yellow-browed Warbler. Autumn. Adult: pattern of wing (11-X) (Photo: Pere Josa)