

378 Western Bonelli's Warbler

Put your logo here



Bonelli's Warbler. Spring (13-IV).



WESTERN BONELLI'S WARBLER (*Phylloscopus bonelli*)

IDENTIFICATION


10-11 cm. Brown grey upperparts, with greenish tinge; white underparts; rump, edges of wing coverts and flight feathers yellow-greenish; white supercilium.



Bonelli's Warbler. Pattern of breast and rump.




Iberian Chiffchaff



Chiffchaff

SIMILAR SPECIES


Iberian Chiffchaff and **Chiffchaff** have dark legs and emargination to 5th primary. **Willow Warbler** has yellow underparts and without limit between rump and mantle. **Wood Warbler** has 10th primary shorter than primary coverts and with contrast between the yellow head and upper breast and pure white underparts. **Melodious Warbler** is bigger, with uniform yellow underparts and broad bill on base.



Willow Warbler

378 Western Bonelli's Warbler

Put your logo here




Wood Warbler



Bonelli's Warbler. Autumn. Ageing. Wear and moulting limit on tail: left adult; right juvenile.



Melodious Warbler



Bonelli's Warbler. Autumn. Ageing. Wear of wing coverts: top adult; bottom juvenile.

SEXING

Plumage of both sexes alike.



AGEING

2 age groups can be recognized:

Juvenile with fresh feathers; yellowish rump with diffuse contrast with mantle.

Adult with worn tail and flight feathers (sometimes some fresh wing covert and flight feathers moulted); yellow rump with distinct contrast with mantle.

In **spring**, after **prebreeding** moult, ageing is not possible using plumage pattern.



Bonelli's Warbler. Autumn. Ageing. Wear and moulting limit on tertials: left adult; right juvenile.

378 Western Bonelli's Warbler

Put your logo here

MOULT

Partial **postbreeding** moult in breeding areas involving body feathers and wing coverts; sometimes also tertials and tail feathers. Partial **postjuvenile** moult involving only body feathers. Both age groups have a complete **prebreeding** moult in wintering quarters (**CAUTION**: birds of both ages may retain some unmoulted flight feathers).

PHENOLOGY

I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XII

STATUS IN ARAGON

Summer visitor. Widely distributed throughout the mountainous forests of the Region.



Bonelli's Warbler. Autumn. Adult (06-VII).



Bonelli's Warbler. Autumn. Juvenile (10-VIII).



Bonelli's Warbler. Head pattern: top in spring (13-IV); middle adult in autumn (06-VII); bottom juvenile (10-VIII).

Bonelli's Warbler. Breast pattern: top left in spring (13-IV); top right adult in autumn (07-VII); left juvenile (10-VIII).

378 Western Bonelli's Warbler

Put your logo here



Bonelli's Warbler. Tail pattern: top left in spring (13-IV); top right adult in autumn (22-VII); left juvenile (10-VIII).



Bonelli's Warbler. Upperpart pattern: top left in spring (13-IV); top right adult in autumn (07-VII); left juvenile (10-VIII).



Bonelli's Warbler. Autumn. Rump pattern: left adult (07-VII); right juvenile (10-VIII).



Bonelli's Warbler. Spring: pattern of wing (13-IV).

378 Western Bonelli's Warbler

Put your logo here



Bonelli's Warbler. Autumn. Adult: pattern of wing (07-VII).



Bonelli's Warbler. Autumn. Juvenile: pattern of wing (10-VIII).